

The species breeds in small numbers in wetlands of the Great Lakes Basin. This basin covers an area of c.106,000km² and extends between approximately 46°-51°N and 90°30'-96°E. The average elevation is about 1,100 m. The basin is divided into three parts: the Uvs, Khyargas and Sharga Lakes Depressions. It is surrounded by mountain ranges of the Altai, the Khangai and the Tagnyn Mountains. The Great Lakes Basin holds a series of lakes that are fed by rivers from mountains of the Altai and the Khangai Mountains. These rivers end in broad deltas and lakes. Some are freshwater lakes, whilst some are saline lakes. The biggest freshwater lakes are the Khar Us Lake and the Khar Lake which are connected by the Chono Kharaiikh Gol. Khar Lake is connected with the Dorgon Lake by a natural channel. The Dzavkhan Gol feeds the freshwater Airag Lake, which drains into the Khyargas Lake. The Buyant Gol and Khovd rivers run from the Altai Mountains and flow into the Khar Us Lake. The Dzavkhan and Khunggui rivers run from the Khangai Mountains and flow into the Airag Lake.

8.7.2 Recent records and distribution

An overview of records of White-headed Duck in Mongolia is summarised in Table 9 and site locations are provided in Map 7.

The White-headed Duck has been recorded in the Kovd Province in the 1990s; 238 birds were recorded in September 1998 at the southern section of Khar Us Lake. There is a report of “a large colony” in June 2000 at Uvs Lake, Uvs Province (D. Batdelger, pers. comm., May 2002).

In recent years, the breeding range of White-headed Duck appears to have extended about 700km to the east, to Olon Lake in the Bulgan Province (N. Tseveenmyadag, pers. comm., April 2002).

8.7.3 Population and trends

Due to a lack of research and monitoring, the population size and trends of White-headed Duck in Mongolia are not clear. Two respondents to the survey provide different population estimates and additional field surveys and information is required.

N. Tseveenmyadag (pers. comm., April 2002) estimates that there are about 500-1000 White-headed Ducks in Mongolia. The approximate number of breeding pairs at the main breeding sites:

- 100 pairs in the Tes River Delta, east of Uvs Lake;
- 5-10 pairs at Shuvuun Tsuglaan Lake, west of Uvs Lake;
- 10-20 pairs in the Zost Lakes, west Airag Lake;
- 100-150 pairs in Khar-Us Lake, Chono-kharaiikh, Khoit Dalai (northern sea), island of White River (Tsagaan Gol); and
- a single observation of 5 birds in central Mongolia in June 2001.

Dashnamjilyn Batdelger (pers. comm., May 2002) believes that the number of White-headed Duck in Mongolia is increasing, and the population is 150-200 individuals. This figure may have been underestimated given a count of 238 birds at Khar Us Lake in September 1998 and the “large colony” at Uvs Lake in June 2000. According to the recent counts, the breeding population of the White-headed Duck in Mongolia could be around 250 pairs.

8.7.4 Conservation status

The White-headed Duck is listed as a rare species in the previous Law on Hunting (1995), the new Law on Fauna (2000) and the Mongolian Red Book (1997).

According to the Mongolian Law on Hunting, Law on Fauna and Red Book, it is prohibited to hunt White-headed Duck. Collection of specimens of White-headed Duck for scientific purposes is permitted under special permission. Mongolians traditionally do not hunt waterbirds. Therefore hunting does not exert much influence on bird numbers.

White-headed Duck are protected under the following Mongolian Laws and regulations: Mongolian Law on Environmental Protection (1995); Mongolian Law on Special Protected Areas (1995); Mongolian Law on Buffer Zones (1997); Mongolian Law on Water (1995); Biodiversity Conservation Action Plan (1996); National Program on Special Protected Areas (1998); National Program on Water (1999); National


Ecological Education Program (1997); Cabinet Ministry Re. 152, Appendix 1, List of Rare Animals (1995).

Table 9: Records of the White-headed Duck in Mongolia, 1980-2002.

Date	No. of birds recorded	Site name and location	Remarks ¹	Counter/source
July 1981	20	Uvs Lake, Uvs Province		D. Batdelger pers. comm., May 2002
End of July 1981	18	Uvs Lake and terminus of Tes River		Boldbaatar Sh. 1997
End of July 1981	8	Khayrgas Lake, Uvs Province		Boldbaatar Sh. 1997
14-17 Aug 1981	15-20/day	Jiree and Tes River Mouth, east side of Uvs Lake	2MA, 2MJ, 3FJ collected	Observation by Sh. Boldbaatar and D. Batdelger. N. Tseveenmyadag, pers. comm., April 2002
19-20 May 1985	3	Shuvuun Tsuglaan Lake, west side of Uvs Lake	1MA	N. Tseveenmyadag, pers. comm., April 2002
5-8 Jun 1985	5-9/day, 20 on last day	Jiree and Tes River Mouth, east side of Uvs Lake	5-9 MA/day, last day about 20AD, 10FA	Observation by N. Tseveenmyadag and D. Batdelger. N. Tseveenmyadag, pers. comm., April 2002
13 Jun 1985	1	Shuvuun Tsuglaan Lake, west Uvs Lake	1MA	Observation by N. Tseveenmyadag and D. Batdelger. Bräunlich and Tseveenmyadag, in prep.
8 May 1995	1	Hodoo Lake, near Zereg, Kovd	1F	Bräunlich 1995
10 May 1995	22	Khar Us Lake (southern section), Kovd	9M,13F	Bräunlich 1995
16 Jun 1995	29	South part of Khar Us Lake (southern section), Kovd	28 M,1 F	Bräunlich 1995
21 Jun 1995	2	Chono Kharaikh River Delta/ Khar Lake, Kovd	1 pair	Bräunlich 1995
13 Jun 1996	12	Khar Us Lake (southern section)	9 M,3F	Observation by M. Köpman. Bräunlich and Tseveenmyadag, in prep.
27 Jun 1998	2	Khar Us Lake	1M,1F	S. Gombobaatar pers. comm., April 2002
24 Sep 1998	238	Khar Us Lake (southern section)	Including c.60 M	Liegl 1998
28-30 Jun 2000	“Large colony”	Uvs Lake		Observation by Valerii Moseikin. D. Batdelger, pers. comm., May 2002
27 Aug 2001	40	Khar Us Lake (southern section)	Minimum number of birds	Observation by C. Bock. Bräunlich and Tseveenmyadag, in prep.
28 Aug 2001	3	Khar Us Lake (southern section)	Minimum number of birds	Observation by C. Bock. Bräunlich and Tseveenmyadag, in prep.
28 Jun 2001	5	Olon Lake, Bulgan	3MA and 2FA	Observation by S. Boldbaatar and G. Mainjagal. N. Tseveenmyadag, pers. comm., April 2002

Note: ¹ M - Male(s), F - Female(s), MA - Male Adult, FA – Female Adult, MJ - Male Juvenile, FJ - Female Juvenile, AD - Adult(s)

Map 7: Distribution of the White-headed Duck in Mongolia during 1980-2002.



No.	Site name	Location
1	Chono Kharaikh River Delta/ Khar Lake	Approx. 48°05'N, 93°10'E
2	Hodoo Lake, near Zereg	Approx. 47°10'N, 92°50'E
3	Khar Us Lake (southern section)	47°45'- 48°23'N, 91°57'- 92°49'E
4	Khayrgas Lake, Uvs Province	48°58'- 49°20'N, 92°48'- 93°48'E
5	Olon Lake, Bulgan	49°52'N, 102°38'E
6	Uvs Lake, Uvs Province	49°59'- 50°41'N, 92°13'- 93°25'E
7	Jiree and Tes rivers Mouth, east side of Uvs Lake	50°25'N, 93°05'E
8	Shuvuun Tsuglaan Lake, west side of Uvs Lake	50°12'N, 92° 10'E

The Uvs Lake Basin was declared a Strictly Protected Area in 1993 and Khar Us Lake and Khyargas Lake were declared as National Parks in 1997 and 2000, respectively. Khar Us Lake was listed as a Ramsar site in 1999.

8.7.5 Threats

Habitat loss and degradation pose significant threats to some of the important sites. Specific threats include reported:

- Construction of a new dam (a planned hydroelectric power station at Chono Kharaikh Gol) will probably destroy breeding sites at Dalai Lake and Khar Lake due to predicted decrease in water levels, increased salinity and decline in aquatic vegetation.
- Livestock grazing on reed beds during winter. Currently reed beds in the southwest corner of Har Us Lake seem to be little affected by grazing during winter. However, a change in this situation might alter or even destroy this nesting area for White-headed Duck (Bräunlich 1995).
- Reed-cutting. At Khar Us Lake and other lakes, where White-headed Duck breed, local people cut the reeds in autumn to build cattle fences for protection of cattle in winter. They drain/dry the wetlands for reed cutting which causes habitat loss for White-headed Ducks.
- Agricultural irrigation activities along the main rivers in dry years.
- Steppe fires that spread into the reed beds and destroy nesting habitat.

- Destruction of reeds by Muskrat that reduce nesting habitat.
- Hunting. Although the species is legally protected, there remains a need for further enforcement of protection and education of local hunters on the legal and conservation status of the species.

8.7.6 Recommendations

The following recommendations have been proposed primarily on the basis of information and feedback from government and non-government respondents to the survey.

- Conduct surveys in potentially suitable wetlands in autumn (August/September) and spring/summer (May/June) to ascertain breeding populations and to identify and confirm the importance of key sites.
- Promote declaration of key sites for White-headed Duck as protected areas, with an emphasis on breeding sites.
- Identify major threats to White-headed Duck including habitat loss and degradation, study the impact or effects of winter grazing of cattle at south Khar Us Lake and incorporate results into a site management plan.
- Undertake a research programme on the White-headed Duck to investigate:
 - characteristics and status of main breeding sites,
 - breeding population size and trends through regular surveying and monitoring,
 - details on breeding biology of the species, including ascertaining clutch size, hatching, fledgling and breeding success and annual mortality,
 - seasonal habitat selection, and
 - feeding requirements and habits.
- Undertake a satellite-tracking project for White-headed Duck to identify the migratory routes and staging sites along the flyway. This will support efforts to develop and implement a comprehensive international conservation action plan.
- Organise a training course on waterbird and wetland conservation for rangers of protected areas.
- Organise an awareness raising and education programme for the local communities living around and within protected areas on the importance to conserve White-headed Duck and other waterbirds.

8.8 PAKISTAN

8.8.1 Historical records and distribution

In Pakistan, the White-headed Duck has been historically recorded in districts of western Pakistan, Punjab, Baluchistan, Bahawalpur (BirdLife International 2001) and Sind (Roberts 1991). However, this species has mainly been recorded in Punjab since the 1960s, at Ucchali, Khabekki, Jahlar, Kallar Kahar, Kharal and Nammal Lakes (see Table 10). In January 1968, a peak count of a total of 1,039 birds was recorded in three wetlands in Punjab. But after 1989, the species has been recorded only from the Salt Range Lakes of Ucchali Complex: Ucchali, Khabekki and Jahlar Lake.

8.8.2 Recent records and distribution

In January 1983 and 1987, there were still 734 and 733 birds counted respectively in Pakistan, but from 1992-1994, only about 150 White-headed Duck recorded. The number of birds rapidly declined after 1995. From 1995-1998, only about 50 birds were recorded every year, and in 2001, only 10 birds were recorded (Chaudhry 2002 and Rahat Jabeen, pers. comm., May 2002).

Peak counts at the Ucchali Complex in January 2002 were very low with different numbers recorded by various observers. A. A. Chaudhry and his colleagues from the Punjab Wildlife Research Institute only

counted 5 birds at the Ucchali Complex, and the number was reduced to 3 in February 2002. However, 10 birds were recorded by Malik Farooq Ahmad, Field Officer of WWF–Pakistan. Zulfiqar Ali, Research Officer of WWF-Pakistan believes there were only 6 birds present in January 2002.

In the latest count carried out by Zulfiqar Ali (pers. comm., November 2002), a total of 34 birds have been recorded on 29 November 2002 at the Ucchali and Jahlar Lakes. At this time, the Khabakki Lake has totally dried up.

An overview of records of White-headed Duck in Pakistan is summarised in Table 11 and site locations are provided in Map 8.

Table 10: Records of the White-headed Duck in Punjab, Pakistan till 1980.

Date	No. of birds recorded	Site name and location	Counter/source
Jan 1967	218	Khabekki Lake	Pakistan Forest Institute Peshawar (PFIP)
Jan 1967	90	Kallar Kahar Lake	PFIP
Jan 1968	1,005	Ucchali Lake	PFIP
Jan 1968	9	Kallar Kahar Lake	PFIP
Jan 1968	25	Nammal Lake	PFIP
Jan 1970	1	Ucchali Lake	PFIP
Jan 1970	1	Khabekki Lake	PFIP
Jan 1970	4	Nammal Lake	PFIP
Jan 1971	161	Khabekki Lake	PFIP
Jan 1971	39	Nammal Lake	PFIP
Jan 1972	102	Ucchali Lake	PFIP
Jan 1972	85	Khabekki Lake	PFIP
Jan 1972	127	Nammal Lake	PFIP
Jan 1973	11	Ucchali Lake	PFIP
Jan 1973	388	Khabekki Lake	PFIP
Jan 1973	111	Nammal Lake	PFIP
Jan 1975	470	Ucchali Lake	PFIP

8.8.3 Population and trends

The White-headed Duck population in Pakistan has greatly decreased since the 1960s, when 1,039 birds recorded in 1968 has reduced to 733 in 1987. In 1994, there were still 148 birds counted in January, however, only about 10 birds were counted in January 2001 and 2002 (Chaudhry 2002). Nevertheless, a total of 34 birds have been recorded on 29 November 2002 at the Ucchali Complex (Zulfiqar Ali, pers. comm., November 2002) but the field staff of WWF-Pakistan believe there are only 14 birds existing at the same time (Rahat Jabeen, pers. comm., November 2002). However it is believed that the numbers of the White-headed Duck is higher than in January 2001 and 2002.

The main reason for this decline in 2000-2002 is considered to be the shortage of rainfall in the area as rainfall has declined in Pakistan over the last five years and the Ucchali wetlands are natural closed basins fed only by rainfall. Some White-headed Duck in the Salt Range may have moved from the Ucchali Complex to surrounding ponds and water reservoirs (Rahat Jabeen, pers. comm., May 2002) where they are not recorded.

8.8.4 Conservation status

A range of measures have been undertaken by government, research institutes and the NGOs in relation to enforcement of legislation, awareness raising and related conservation measures.

Legislation: the White-headed Duck is legally protected in all provinces and federal units in Pakistan. It is included on Schedule 3 of protected animals under the Punjab Wildlife Protection, Conservation and Management Act 1974.

Table 11: Records of the White-headed Duck in Punjab, Pakistan, 1980-2002.

Date	No. of birds recorded	Site name and location	Counter/source
Jan 1983	734	Khabekki Lake	Pakistan Forest Institute Peshawar (PFIP)
Jan 1984	209	Khabekki Lake	PFIP
Jan 1984	46	Kallar Kahar Lake	PFIP
Jan 1985	120	Khabekki Lake	PFIP
Jan 1986	325	Khabekki Lake	PFIP
Jan 1987	620	Ucchali Lake	Punjab Wildlife Research Institute (PWRI)
Jan 1987	65	Jahlar Lake	PWRI
Jan 1987	48	Kharal Lake	PWRI
Jan 1988	154	Ucchali Lake	PWRI
Jan 1988	132	Jahlar Lake	PWRI
Jan 1989	100	Ucchali Lake	PWRI
Jan 1989	56	Jahlar Lake	PWRI
Jan 1989	31	Kharal Lake	PWRI
Jan 1990	18	Ucchali Lake	PWRI
Jan 1990	44	Jahlar Lake	PWRI
Jan 1990	14	Kharal Lake	PWRI
Jan 1991	2	Ucchali Lake	PWRI
Jan 1991	32	Khabekki Lake	PWRI
Jan 1991	30	Jahlar Lake	PWRI
Jan 1992	144	Khabekki Lake	PWRI
Jan 1992	2	Jahlar Lake	PWRI
Sep 1992	2	Jahlar Lake	PWRI
Oct 1992	12	Jahlar Lake	PWRI
Nov 1992	145	Jahlar Lake	PWRI
Dec 1992	50	Khabekki Lake	PWRI
Dec 1992	81	Jahlar Lake	PWRI
Jan 1993	36	Khabekki Lake	PWRI
Jan 1993	78	Jahlar Lake	PWRI
Feb 1993	43	Khabekki Lake	PWRI
Feb 1993	68	Jahlar Lake	PWRI
Mar 1993	43	Khabekki Lake	PWRI
Mar 1993	18	Jahlar Lake	PWRI
Apr 1993	3	Jahlar Lake	PWRI
May 1993	2	Jahlar Lake	PWRI
Jun 1993	2	Jahlar Lake	PWRI
Jul 1993	3	Jahlar Lake	PWRI
Aug 1993	3	Jahlar Lake	PWRI
Nov 1993	25	Khabekki Lake	PWRI
Nov 1993	111	Jahlar Lake	PWRI
Dec 1993	148	Jahlar Lake	PWRI
Jan 1994	144	Jahlar Lake	PWRI
Feb 1994	121	Jahlar Lake	PWRI
Mar 1994	43	Jahlar Lake	PWRI
Apr 1994	94	Jahlar Lake	PWRI
Jan 1995	14	Khabekki Lake	PWRI
Jan 1995	37	Jahlar Lake	PWRI
Jan 1996	32	Jahlar Lake	PWRI
Jan 1997	36	Khabekki Lake	PWRI
Jan 1997	16	Jahlar Lake	PWRI
Jan 1998	1	Ucchali Lake	PWRI
Jan 1998	3	Khabekki Lake	PWRI
Jan 1998	9	Jahlar Lake	PWRI
23 Mar 1998	9	Khabekki Lake	Zulfiqar Ali
23 Mar 1998	47	Jahlar Lake	Zulfiqar Ali

Date	No. of birds recorded	Site name and location	Counter/source
Oct 1998-Mar 1999	9	Khabekki Lake	Zulfiqar Ali
Oct 1998-Mar 1999	27	Jahlar Lake	Zulfiqar Ali
Jan 1999	8	Khabekki Lake	PWRI
21-Feb 1999	12	Jahlar Lake	PWRI
Mar 1999	8	Khabekki Lake	Malik Farooq Ahmad
Oct 1999	6	Khabekki Lake	Malik Farooq Ahmad
Nov 1999	6	Khabekki Lake	Malik Farooq Ahmad
Jan 2000	4	Khabekki Lake	PWRI
03-Jan 2000	7	Khabekki Lake	Zulfiqar Ali
03-Jan 2000	16	Jahlar Lake	Zulfiqar Ali
Feb 2000	6	Jahlar Lake	Malik Farooq Ahmad
Jan 2001	5	Khabekki Lake	PWRI
Jan 2001	10	Jahlar Lake	Zulfiqar Ali, Malik Farooq Ahmad
Jan 2002	10	Jahlar Lake	Malik Farooq Ahmad
Jan 2002	6	Jahlar Lake	Zulfiqar Ali
Jan 2002	5	Jahlar Lake	PWRI
Feb 2002	3	Jahlar Lake	PWRI
21 Oct 2002	9	Uchalli Lake	Zulfiqar Ali
21 Oct 2002	18	Jahlar Lake	Zulfiqar Ali
15 Nov 2002	11	Uchalli Lake	Zulfiqar Ali
15 Nov 2002	18	Jahlar Lake	Zulfiqar Ali
Nov 2002	14	Ucchali Complex	WWF-Pakistan
29 Nov 2002	14	Uchalli Lake	Zulfiqar Ali
29 Nov 2002	20	Jahlar Lake	Zulfiqar Ali
8 Dec 2002	11	Uchalli Lake	Zulfiqar Ali
8 Dec 2002	15	Jahlar Lake	Zulfiqar Ali

Protected Areas: Khabekki, Jahlar, Nammal and Kharal have all been declared as Wildlife Sanctuaries where all habitat disturbance, hunting and trapping is prohibited. Uchchali Lake has been declared a Game Reserve where hunting is allowed only on a special permit, but such permits have never been granted. The Uchchali Complex, which includes Khabekki, Uchchali and Jahlar Lakes was designated as a Ramsar site in March 1996.

Management: A management plan was formulated by WWF and Punjab Wildlife and Parks Department in 1994. The plan was revised subsequently by the Department in 1999.

The Government of Pakistan is currently developing a GEF/UNDP funded project for “Conservation of wetlands in Pakistan” wherein four wetlands are to be taken up for management planning and subsequent implementation.

Awareness/Education: An awareness campaign has been conducted in the area by the Punjab Wildlife and Parks Department since the late 1980s. WWF-Pakistan partly funded a project entitled “Conservation of Wetlands” during 1992 to 1996. Participatory Rural Appraisal exercises were conducted jointly by WWF-Pakistan and the Punjab Wildlife and Parks Department to gauge the community participatory potential in the area in December 1994. The Soan Valley Biodiversity Conservation Forum (formed by WWF-Pakistan and Soan Valley Development Programme) are also actively working in the area to help educate the public on conservation measures to be adopted to save the biodiversity/natural resources of the area. WWF-Pakistan is also running an awareness campaign focusing their activities on the conservation of waterbirds and wetlands.

The White-headed Duck has been proposed for inclusion in the list of threatened bird species (Red Data Book) currently being compiled by WWF-Pakistan. Within the South Asian region, Pakistan host almost the entire wintering population of the White-headed Duck and therefore, the country has a special responsibility for the conservation of the species (Roberts 1991). The recent drastic decline in its wintering population in Pakistan warrants immediate action.

Map 8: Distribution of the White-headed Duck in Pakistan during 1980-2002.



No.	Site name	Location
1	Jahlar Lake	32°29'N, 72°07'E
2	Kallar Kahar Lake	32°46'N 72°42'E
3	Khabekki Lake	32°37'N 72°14'E
4	Kharal Lake	30°53'N, 73°35'E
5	Ucchali Lake	32°33'N, 72°01'E

8.8.5 Threats

Threats to the wintering population of White-headed Duck are mainly related to habitat loss and modification, competition with fisheries, and to a lesser extent, hunting and disturbance.

- **Habitat Loss and Modification:** The main threat to the White-headed Duck population in Pakistan is habitat degradation and loss. The shallow wetlands occupied by the duck fluctuate naturally with climatic conditions such as rainfall. The intensity of rainfall and the amount of run-off determine the well being of the wetlands. Failure of rainfall in recent years has adversely affected wetlands and their extent has greatly been reduced. All three wetlands in the Ucchali Complex are surrounded by agricultural fields. Fertilisers and pesticides are frequently used. Leaching and run-off of fertilisers and pesticides in surface water pollutes the wetlands, although their impact has not been determined. Ucchali Lake is particularly prone to this source of pollution. The off-season cauliflower *Brassica* sp. crop, which takes only six weeks to mature, is the main cash crop in Ugali village on the northern edge of the lake. To increase yields, fertiliser and pesticide use is quite common. This uncontrolled usage results in the pollution of the lake. Over-abstraction of groundwater, both for drinking and for agricultural purposes, also causes a lowering of the water table and a subsequent reduction in the extent of lakes/wetlands. Most abstraction takes place in the Ucchali village on the southern edge of the Ucchali Lake.

The land around the wetlands is privately owned and any reduction in the extent of the lakes prompts landowners to start cultivating exposed areas. This practice is most destructive at Khabekki Lake where the owners have cultivated the land right up to the edge of the water. At Ucchali Lake, the water has receded, but as the land has turned into a marsh, it is not suitable for cultivation. Noxious

gases emanating from the marsh make the habitat unsuitable for tourists. Jahlar is the only wetland that has so far retained its character to some extent as the exposed lakebed has not been cultivated.

In summary, wetland habitat has been lost at Khabekki Lake, degraded at Ucchali Lake and reduced at Jahlar Lake.

- **Overgrazing:** Vegetated areas around the lakes are heavily grazed by domestic livestock - buffaloes, cattle, sheep, goats, donkeys and camels. Grazing is much beyond the grazing capacity levels as found in the Participatory Rural Assessment exercise undertaken by WWF-Pakistan and Punjab Wildlife and Parks Department (1995).
- **Fishing and fish introductions:** Fishing activities at Khabekki threaten waterbirds in a number of ways: introduced fish compete for food with waterbirds, birds are caught and drown in fishing nets and fishing activities disturb waterbirds. White-headed Duck, however, are known to leave the lake during such periods when they are thought to disperse to Jahlar Lake.

The population expansion of the introduced *Tilapia Oreochromis* sp. in Khabekki Lake resulted in eutrophication and mortality. These fishes were exterminated by the Fisheries Department after which various kinds of carp were introduced. These fish survived until they were killed by drought conditions in 2001. Fisheries operations have since ceased at the lake.

Ucchali Lake has been considered unsuitable for the introduction of fish due to the pH and salinity of the lake. Jahlar is thought to be too small for fishing operations to be economically viable.

- **Hunting and Disturbance:** Hunting is not allowed on any of the aforementioned lakes. Instances of illegal hunting and poaching have been reported but not in recent years. White-headed Ducks could, however, be shot by mistake by hunters who are unable to identify the species.
- **Recreation and disturbance:** Recreational activities at Kallar Kahar; and boating operations at Ucchali Lake cause disturbance and affect waterbird populations.

Three other lakes, two in the Salt range –Nammal and Kallar Kahar, and the third (Kharal) in the central Punjab Plains, have already lost their habitat characteristics due to human interference and excessive recreational activities. The White-headed Duck no longer visits these lakes. At Nammal, fish have been introduced and the lake water is now being used for irrigating agricultural fields downstream. Kallar Kahar has now been developed into a recreational resort and due to disturbance, very few waterbirds visit the lake. Kharal was developed as a drainage basin to remove water from surrounding agricultural fields. Tilapia and carp were introduced into the lake. In the early 1990s reclamation projects were then started and the excess water was drained from the agricultural fields all around the lake hence the inflow of water to the lake was stopped resulting into the deterioration of habitat characteristics. The human population surrounding the wetland has increased and as a result of that, reclamation of land has increased.

8.8.6 Recommendations

The following recommendations have been proposed primarily on the basis of information and feedback from government and non-government respondents to the survey.

- Undertake a study of the status of the wintering population of White-headed Duck on the Salt Range Lakes in northern Pakistan. This should be undertaken in relation to its status on the breeding grounds in Central Asian States to determine the causes of its population decline, and to identify remedial measures for its conservation.
- Institute long term monitoring studies of climatic factors and their effect on the physical, chemical and biological characteristics of the Salt Range Lakes.
- Undertake a study on the status of the White-headed Duck on the Salt Range Lakes in relation to changes in physical, chemical and biological characteristics of the lakes to suggest appropriate habitat development to ensure the survival of the species.

- Undertake a satellite-tracking project to identify the migratory route and stopover sites along the flyway of the White-headed Duck.
 - Strengthen staff capacity of protection agencies at the Salt Range Lakes to prevent illegal hunting of waterbirds which could lead to further population decline.
 - Improve habitat conditions at the Salt Range Lakes, for example, afforestation in surrounding watersheds and restoration of water resources prevent silting of the lakes and further habitat loss.
 - Provide training to local communities on efficient use of local fuels to minimise the use of vegetation cover at the Salt Range Lakes.
 - Undertake public awareness raising programmes for villages around the Salt Range Lakes through organising local Community Based Organisations/Village Organisations (CBOs/VOs) to increase conservation awareness.
 - Encourage development of sustainable ecotourism at the Salt Range Lakes to help improve the local economy.
 - Ensure management of the Salt Range Lakes, on a Participatory/Benefit sharing basis, with the active involvement of local communities.
-

8.9 RUSSIA (Asian part only)

8.9.1 Historical records and distribution

In the Russian Federation, the White-headed Duck is largely a breeding species of the lowland forest-steppe wetlands and is mainly recorded during the summer months. It occurs during spring and autumn months on migration across through the southwest of the country.

An overview of records of White-headed Duck in Asian part of Russia is summarised in Table 12. Breeding occurs to the east of the Urals in the Chelyabinsk Oblasts on the border of west Siberia and Kazakhstan, (BirdLife International 2001, Gordienko 1986). Breeding also takes place on the wetlands of the Tobol-Ishim Forest-steppe in the eastern part of Kurgan Oblast and in the southern part of the Tyumen Oblast in West Siberia (Krivenko 1999, Linkov 2001). However, the most important breeding area is now thought to be further east along the border, in the Kulunda and Baraba forest-steppes in Novosibirsk Oblast (BirdLife International 2001). The White-headed Duck is recorded as an occasional breeding species at the Chany Lakes and in the Lower Bagan area, 100-150 pairs were recorded in 1970 and 1971 (Krivenko 1999). Further eastwards it has been recorded breeding on Ubsu-Nur Lake in South Tuva on the Mongolian border (Zabelin *et al.* 2000) and it also occurred on the Tore-Khol Lake of this area (Krivenko 2000).

In the European part of Russia, breeding used to occur in the Sarpa lowlands between Volgograd and the Caspian and in the Volga/Ural Steppes. The species has also been recorded in the northern Caucasus and along the western coast of the Caspian, while the Manych Valley is a major spring and autumn migration site for the species (Green and Anstey 1992).

In Daghestan according to data gleaned from hunters, the White-headed Duck appears sometimes on the Caspian Sea and coastal water bodies in November-December. In the years from 1970 to 1980 in the central Caspian shore during winter, up to 300 birds have been recorded (Red Data Book of Republic of Daghestan 1998).

8.9.2 Recent records and distribution

Recently, White-headed Duck have been recorded breeding in many large waterbodies of the forest-steppe part of the Ural-Chelyabinsk Region. This includes Momynkul Lake, Zabaluevo Lake, Katai Lake,

Travyanoe Lake, Kurlady Lake and the lake near Cheraskul (Gordienko *et al.* 1986, Gordienko 1998, Gordienko 2001, Karyakin and Kozlov 1999, Karyakin 1998, Braude 1989). During 1992-1996, Karyakin *et al.* (1999) observed a total of 298 White-headed Duck (including ducklings) and 48 broods on 29 lakes of the forest-steppe Trans-Uralia. They calculated a breeding density of 2-7 pairs per 1km² in nest-suitable habitats and 2-10 pairs per 10km² over the total area. The main breeding area of the species in the Chelyabinsk region is situated within the border of the forest-steppe Trans-Uralia between 54°15'-56°05'N, 58°57'-61°30'E.

In the Sverdlovsk Region, the White-headed Duck is a rare vagrant. Braude (1989) gives information, based on the word of hunters, about sighting of the species near Nikol'skoe settlement (southern border of the Sverdlovsk Region). Karyakin *et al.* (1999) observed White-headed Duck at Chervyanoe Lake on 15 May 1995.

In the Omsk Region, 4 birds were sighted in the Stepnoi (steppe) Game Reserve: Ataich'ie Lake on 15 May 1997 in a mixed flock of Tufted Ducks *Aythya fuligula* and Pochards *Aythya ferina*. Furthermore, two pairs were spotted at Sylkino Lake at the end of May 1998 (Yakimenko 1998) and 7 birds on 21 May 2000 (Yakimenko V. V. unpublished).

In the Novosibirsk Region, White-headed Duck have been recorded breeding at Chany and Baganskiy Lakes during the summer of 1969-1989 (Gordienko *et al.* 1986). A. K. Yurlov (pers. comm., May 2002) has recorded White-headed Duck breeding in the summer in the Chany Lake, Fadiha Lake, Ulianovskoe Lake and Orlovka Village Lake since 1997.

The species was also recorded in Khakassia, to the southwest of the Krasnoyarsk Region, on Shira Lake and Belye Lake in the 1980s (Syroechkovski and Rogacheva 1995).

In the European part of Russia, in the Orenburg Region, 30 White-headed Duck were shot during the spring passage in the 1980s (Gordienko 1998) and one female was bagged by a hunter on 30 August 1999 at the sewage reservoir of the Svetlyi settlement (Korshikov and Kornev 1999).

On 27 October 1980, 1,200 White-headed Duck were counted on a 100km transect bisecting the Manych Lake and Manych-Gudilo Lake. Several pairs also occasionally breed in this area (Linkov 1984, Green and Anstey 1992).

Table 12: Records of the White-headed Duck in Asian part of Russia till 1980.

Date	No. of birds recorded	Site name and location	Remarks	Counter/source
15 May 1940	2	Chebarkul Lake, Ural (55°07'N, 60°35'E)	migrating	Observation by Migun N.N. (Gordienko N. S. pers. comm, April 2002)
May-Sep 1969-1989	230 pairs	Chany and Baganskiy Lakes, southwest Siberia, Russia. Presnovskiy and Mibalykskiy Lakes and Naurzumskiy Lakes in northern Kazakhstan	migrating and nesting ¹	Gordienko <i>et al.</i> 1986 Gordienko 1995, 1998
18 June 1969	1	Butash Lake, Chelyabinsk Region	male	Karyakin and Kozlov 1999
13 May 1972	2	Katai Lake, Ural	migrating	Gordienko 1998
1979	Female with chicks	Kulundinskoe Lake, Altaj krai, Novosibirsk Region (52°57'N, 79°43'E)		Kuchin 1991
23-25 August 1979	Flock	Gor'koe Lake, Altaj Kari, Novosibirsk Region	moulting males	Irisov and Irisova 1982

Note: ¹The record refers to southern Russia and northern Kazakhstan

Table 13: Records of the White-headed Duck in Asian part of Russia, 1980-2002.

Date	No. of birds recorded	Site name and location	Remarks	Counter/source
May-Sep 1969-1989	230 pairs	Chany and Baganskiy Lakes, southwest Siberia, Russia. Presnovskiy and Mibalykskiy Lakes and Naurzumskiy Lakes in northern Kazakhstan	The record refers to southern Russia and northern Kazakhstan	Gordienko <i>et al.</i> 1986 Gordienko 1995, 1998
June 1980-2001	20	Kurlady Lake, Ural, Chelyabinsk Region	Nesting each year	Gordienko 2001
1980s	1	Travyanoe Lake, Chelyabinsk Region		Gordienko 1998
May 1980	5	Shira Lake, Krasnoyarsk Region		Rogacheva 1992
July-Aug 1985	20	Belye Lake, Krasnoyarsk Region		Syroechkovski and Rogacheva 1995
24 July 1987	2	Lake nr. Cheraskul , Chelyabinsk Region	Duckling	Braude 1989
Early Sep 1987	1	Momynkul Lake, Chelyabinsk Region	non-flying brood	Braude 1989
June 1990-1995	6	Zabaluevo Lake, Ural, Oktyabrskiy district, Chelyabinsk Region	nesting each year	Gordienko 2001
End May 1992	3 nests	Kurlady Lake, Chelyabinsk Region	birds with duckling	Karyakin and Kozlov 1999
July 1992	11	Kurlady Lake	Broods	Karyakin and Kozlov 1999
July 1992	7	Sykandyk Lake, Chelyabinsk Region	Broods	Karyakin and Kozlov 1999
July 1992	1	Katai Lake, Ural, Chelyabinsk Region	Brood	Karyakin and Kozlov 1999
July 1992	4	Treustan Lake, Chelyabinsk Region	Broods	Karyakin and Kozlov 1999
July 1992	2	Atkul Lake, Chelyabinsk Region	Broods	Karyakin and Kozlov 1999
July 1992	3	Selezyan Lake, Chelyabinsk Region	Broods	Karyakin and Kozlov 1999
July 1993	3	Tishki Lake, Chelyabinsk Region	Broods	Karyakin and Kozlov 1999
End May 1994	3 nests	Mayan and Alakul Lakes, Chelyabinsk Region		Karyakin and Kozlov 1999
July 1994	2	Dengino Lake, Chelyabinsk Region	Broods	Karyakin and Kozlov 1999
July 1994	1	Utich'ie Lake, Chelyabinsk Region	Broods	Karyakin and Kozlov 1999
July 1994	1	Kartabyz Lake, Chelyabinsk Region	Broods	Karyakin and Kozlov 1999
July 1994	3	Butash Lake, Chelyabinsk Region	Broods	Karyakin and Kozlov 1999
July 1994	5	Malyi Sarykul Lake, Chelyabinsk Region	Broods	Karyakin and Kozlov 1999
July 1994	2	Duvankul Lake, Chelyabinsk Region	Broods	Karyakin and Kozlov 1999
15 May 1995	Observed	Chervyanoe Lake, Sverdlovsk Region		Karyakin <i>et al.</i> 1999
27 May 1995	2 nests	Uelgi Lake, Chelyabinsk Region	birds with duckling	Karyakin and Kozlov 1999
27 May 1995	1 nest	M. Bugodak Lake, Chelyabinsk Region	birds with duckling	Karyakin and Kozlov 1999
1995 and 1997	Recorded	Chebarkul Lake, Chelyabinsk Region		Karyakin and Kozlov 1999; Karyakin 1998

Date	No. of birds recorded	Site name and location	Remarks	Counter/source
19 June 1996	1 nest	Tishki Lake	birds with duckling	Karyakin and Kozlov 1999
July 1996	2	Dengino Lake	Broods	Karyakin and Kozlov 1999
15 May 1997	4	Ataich'ie Lake, Omsk Region	2M+2F ¹	Yakimenko 1998
June 1997	1F	Chany Lake, Novosibirsk Region	non breeding	Yurlov A.K. and G.C.Boere, unpublished
Sep 1997	1M	Fadiha Lake, Zdvinskij district, Novosibirsk Region	killed by hunter	Yurlov A.K. unpublished
End May 1998	2 pairs	Sylkino Lake, Omsk Region		Yakimenko 1998
16 Sep 1998	10 (4F+6 chicks)	Uljanovskoe Lake, Baganskij district, Novosibirsk Region	hunting inspectors have also observed nesting birds at the site in 1995-96	Yurlov A.K. unpublished
21 May 2000	7 (4F+ 3M)	May-Sor Lake, Omsk Region	breeding behaviour observed	Yakimenko V. V. unpublished
5 June 2000	1M	Orlovka village Lake, Chistoozernij district, Novosibirsk Region	breeding behaviour observed	Yurlov A.K. unpublished

Note: ¹M - Male(s), F - Female(s)

During the summers of the 1987-1991, a few White-headed Duck were observed every year in the Vetelka settlement of the Saratov Region (Moseikin 2000). On the reservoirs on Sarpa lowland near Volgograd City, the total breeding number of the White-headed Duck in 1999-2001 is estimated at 75-100 pairs (Bukreev and Chernobay 2002). These ducks were also observed in small numbers during summer in Tsagan-Noor Lake (in the Kalmykia part of Sarpa Lowland) (Shubin 2001).

The current number on migration in Daghestan does not exceed 100 birds, 2 cases of White-headed Duck shootings are known from the Tarum and Derbent Districts (Jamirzoev, 1999). In 2001, the White-headed Duck is found breeding (8-11 pairs) in Adzhi Lake in Daghestan (Dzhamirzoev 2002).

An overview of records of White-headed Duck in Asian part of Russia is summarised in Table 13 and site locations are provided in Map 9.

8.9.3 Population and trends

Asian Part of Russia - A. K. Yurlov (pers. comm., June 2002) has suggested the decreasing trends of the White-headed Duck in the Baraba forest-steppe and Kulunda steppe, West Siberia. During the mid-1970s, there were about 80-150 pairs compared to 30-40 pairs today. The main cause has been recent drought conditions and a subsequent decrease in water levels in the lakes. Two long droughts have been experienced during the last 30 years in southwestern Siberia. However, there has also been an increase in human disturbance in many lakes (e.g. fishing, hunting and other activities).

In the other territories of the Asian part of Russia, Linkov (2001) has estimated the White-headed Duck population to be as follows:

- Tobol-Ishim forest steppe, and Chelyabinsk Region – a few pairs. Krivenko (1999) estimated that there were 5-50 pairs in this region whilst Gordienko (pers. comm., May 2002) suggested a total of 30 pairs in the Southern Ural in the 1990s.
- Tyumen region: 20-30 pairs.
- Khakassia and Tuva: 40-50 pairs

Map 9: Distribution of the White-headed Duck in the Asian part of Russia during 1980-2002.



No.	Site name	Location
1	Ataich'ie Lake, Omsk Region	54°30'N, 75°40'E
2	Baganskiy Lakes, southwest Siberia	54°09'N, 78°23'E
3	Belye Lake, Krasnoyarsk Region	54°40'N, 90°10'E
4	Chany Lakes, southwest Siberia	54°52'N, 77°27'E
5	Chebarkui Lake of Ural, Chelyabinsk Region	55°07'N, 60°35'E
6	Duvankul Lake, Chelyabinsk Region	54°40'N, 61°30'E
7	Fadiha Lake, Zdvinskij district, Novosibirsk Region	54°36'N, 78°12'E
8	Kartabyz Lake, Chelyabinsk Region	54°30'N, 62°30'E
9	Katai Lake, Ural, Chelyabinsk Region	55°04'N, 62°05'E
10	Kurlady Lake, Ural, Chelyabinsk Region	55°01'N, 61°05'E
11	Malyi Sarykul Lake, Chelyabinsk Region	54°50'N, 61°30'E
12	May-Sor Lake, Omsk Region	54°27'N, 75°38'E
13	Orlovka village Lake, Chistoozernij district, Novosibirsk Region	54°28'N, 76°39'E
14	Shira Lake, Krasnoyarsk Region	54°30'N, 90°10'E
15	Tishki Lake, Chelyabinsk Region	55°50'N, 61°40'E
16	Uelgi Lake, Chelyabinsk Region	55°40'N, 61°30'E
17	Ulianovskoe Lake, Baganskij district, Novosibirsk Region	54°08'N, 78°11'E
18	Zabaluevo Lake, Ural, Chelyabinsk Region	54°06'N, 63°00'E

European part of Russia

- Krivenko (2000) estimated 17-20 pairs on the lakes of the Sarpa lowland near Volgograd City. Bukreev and Chernobay (2002) believe there are 75-100 pairs from 1999 to 2001.
- Adzhi Lake in Dagestan, 8-11 pairs in 2001 (Dzhamirzoev 2002).

Linkov (2001) estimated that the total Russian breeding population of White-headed Duck was 170-230 pairs. However, Sergey Bukreev (pers. comm., October 2002) suggests the current estimate can be increased to a minimum of up to 300-500 pairs.

8.9.4 Conservation status

The species is included in the national Red Data Book (2001) as Endangered – 1 category (close to Extinction) and provincial (Chelybinskiy) Red List (in press) as threatened. Wildlife Law (1995) is concerned with protection of rare species in Russia (Linkov 2001). Hunting of White-headed Duck is illegal in Russia.

Some of most important sites for White-headed Duck are protected, mainly as non-hunting areas or "Zakazniks" (corresponding to IUCN category IV for protected areas). White-headed Duck occur in a total of 9 protected areas (with different protected status) in the Asian part of Russia and 5 in the European part.

Regular monitoring of summer numbers and distribution of White-headed Duck in the Chelyabinsk region and European part at Volgograd and Daghestan Regions are being conducted.

8.9.5 Threats

Poaching, particularly in Russia, seems not to be a significant problem. The population is declining perhaps due to habitat loss caused by river flow control and the natural cyclical decrease of steppe wetlands. The other limiting factor is that only a few birds participate in the breeding while most adult individuals remain non-breeding (Linkov 2001).

8.9.6 Recommendations

The following recommendations have been proposed primarily on the basis of information and feedback from government and non-government respondents to the survey.

- Undertake a survey and monitoring programme on the breeding grounds of White-headed Duck. This should include field work for checking of the previous breeding places in the Kurgan, Tyumen, Novosibirsk, Tuva and Khakassia Regions.
- Undertake a satellite-tracking project to identify migratory routes and stopover sites along the flyway of the White-headed Duck. The results will support efforts to develop and implement a comprehensive international species conservation action plan.
- Undertake a ringing (banding) project for the White-headed Duck to identify migratory routes and wintering sites in the flyway.
- Produce and disseminate relevant awareness raising materials targeted mainly at hunters on the need to conserve White-headed Duck and other threatened waterbirds.
- Provide training for game managers and rangers to enable them to identify White-headed Duck and to prevent from illegal capture.

8.10 TAJIKISTAN

8.10.1 Historical records and distribution

In the beginning of the 20th century, White-headed Duck were recorded during the breeding season and on migration on lakes of Vakhsh River Valley (approximately 37°15'N, 68°50'E), southwestern Tajikistan, and in Syr Darya River Valley (approximately 40°20'N, 70°00'E), northern Tajikistan (Ivanov 1940, Ivanov 1969, Isakov 1952, Abdusalamov 1971).

8.10.2 Recent records and distribution

Over the last 30 years there has not been a single record about this species in Tajikistan. However Isom Abdusalamov (1971 and pers. comm., September 2002) considers that the White-headed Duck is a very rare species in Tajikistan and may occur in small numbers on freshwater and brackish lakes with reed brakes and open stretches. It may occur in the Syr Darya and Zerafshan Valleys of northern Tajikistan and in Vakhsh, Pyanj and Kafirnigan river basins of southwestern Tajikistan.

8.10.3 Population and trends

The population of breeding pairs is likely to be decreasing because most former habitats have been developed for agriculture and therefore have become unsuitable for White-headed Duck.

8.10.4 Conservation status

Due to the unclear status of the White-headed Duck in Tajikistan, it was not included in the Red Data Book of Tajikistan (1988).

The accession of Tajikistan to the Convention on Wetlands in 2002 offers opportunities of greater international collaboration and resource mobilisation to undertake wetland and waterbird related conservation work.

8.10.5 Threats

Agricultural development in wetlands and hunting of waterbirds are reported as a general threat in many parts of the country.

8.10.6 Recommendations

The following recommendations have been proposed primarily on the basis of information and feedback from government and non-government respondents to the survey.

- Conduct survey activities in the southwest and northern parts of the country to improve the understanding of the status and distribution of White-headed Duck.
- Protect key sites for White-headed Duck from hunting and other disturbances and changes in ecological character of the wetlands.

8.11 TURKMENISTAN

8.11.1 Historical records and distribution

The White-headed Duck is a common wintering species and passage migrant in Turkmenistan. Historically, birds normally winter and migrate through the southeastern part of the country, along the coast of the Caspian Sea and nearby inland lakes. In February 1932, 47,080 birds were recorded in the Southeast coast of the Caspian Sea (Laptev *et al.* 1934) (See Table 14). The White-headed Duck was also common in these sites in the winter of 1935-1939, and groups of 400-500 birds were regularly observed in Krasnovodsky Bay during southward migration (Isakov and Vorobyev 1940). Other wintering sites have been recorded along the valleys of Amu Darya, Tezhen and Murgab Rivers since at least the 1950s. The peak count of White-headed Duck during the mid-winter waterbird census for the region was 850 in 1974, with 600 at Kelif Lake (Green and Anstey 1992).

Southward migration along the coast of Turkmenistan and the Western Uzboy River Valley starts in October. Key staging sites include the Krasnovodsky and Severo-Chelekensky and Becovich Bays. Northward migration takes place mainly between 11 February to 18 March (Vasilyev and Gauser 2001).

Dementiev and Gladkov (1952) mentioned records of White-headed Duck breeding along the shores of the Lower Amu Darya and Tedzhen/Murgab rivers.

8.11.2 Recent Records and distribution

The most recent surveys for the species were conducted by M. E. Gauser and V. I. Vasilyev, and the results are provided below. From 1986 onwards, most birds have been found at Krasnovodsky and Severo-Chelekensky Bays (86-100% of birds). A total of 820 White-headed Duck were counted in January 1998 along the coast of the southeastern Caspian between Carabogasgol and Gasankuly, where 723 birds were also counted in November 2001 (Vasilyev and Gauser 2001).

Table 14: Records of the White-headed Duck in Turkmenistan till 1980.

Date	No. of birds recorded	Site name and location	Remarks ¹	Counter/source
Feb 1932	47,080	Southeast coast of Caspian Sea		Laptev <i>et al.</i> 1934
Jan 1937	500	Krasnovodsky Bay, Caspian Sea		Isakov and Vorobjev 1940
Jan 1973	490	Mainland lakes/rivers		Poslavski 1992
Jan 1974	850	Mainland lakes/rivers	600 at Kelif Lake	Poslavski 1992
Jan 1974	About 200	Reservoir in the Kara-Kum desert		Vasilyev <i>et al.</i> 1984
25 Sep 1975	44	Krasnovodsky Bay, Caspian Sea	groups of 3-5 birds and individuals	Vasilyev <i>et al.</i> 1984
Jan 1976	373	Lakes/rivers of mainland Turkmenistan		Poslavski 1992
Jan 1977	6	Maloye Delili Lake in the lower Atrek (37°30'N, 54°28'E)		Poslavski 1992
10 Nov 1977	208	Severo-Chelecensky Bay, Caspian Sea	4 flocks: 1M+F, 2-4M+14F, 3-5F, 4-30% male	Vasilyev <i>et al.</i> 1984
Jan 1978	22	Mainland lakes/rivers		Poslavski 1992
11 Jan 1978	73	Krasnovodsky and Severo-Chelecensky Bay, Caspian Sea	13 flocks (1-19 birds) and individual birds	V.I.Vasilyev count. Ataev <i>et al.</i> 1978
Jan 1979	21	Mainland lakes/rivers		Poslavski 1992
11 Feb 1979	82-flight	Krasnovodsky and Severo-Chelecensky Bay	2 flocks	Vasilyev and Gauser 2001

Note: ¹M - Male(s), F - Female(s)

Breeding has been formerly recorded along the middle Amu Darya at Soltantagt Lake in eastern Turkmenistan during 1984-1991, where 5-6 broods of ducklings were seen in May 1987 and 19 breeding pairs were recorded in 1989. However, no birds have been recorded in this area in winter (Poslavski 1992, Green 1992). Further more, M. E. Gauser (pers. comm., May 2002) has also recorded a pair of White-headed Duck nesting at the Krasnovodsky Bay of the Caspian Sea during 21-22 May 1982 and 8 birds in the area in April 2002.

An overview of records of White-headed Duck in Turkmenistan is summarised in Table 15 and site locations are provided in Map 10.

8.11.3 Population and trends

In the latter half of the 20th century, White-headed Duck numbers have fallen from a maximum of 47,080 birds recorded in February 1932. In autumn-winter 1972-1978, records of peak annual counts were 170-600 birds (Ataev *et al.* 1978), while from 1988 onwards, peak counts were down to 19-820 birds (Vasilyev and Gauser 2001). No conclusion can be drawn on the trend or status of breeding populations at this time due to inadequate information (M. E. Gauser and V. I. Vasilyev, pers. comm., May 2002).

8.11.4 Conservation status

The White-headed Duck is listed as an uncommon species in the second edition of the national Red Data Book (1999).


Legislation and regulations relating to White-headed Duck in Turkmenistan include: Act about preservation and rational usage of fauna (1997); Act about Protected Areas (1992); The Model Statute about "Governmental Nature Reserves of Turkmenistan" (1994); The Model Statute about Governmental Arboretums of rare and threatened animals and plants in Turkmenistan (1995); Completion of a "National Action Plan on Biodiversity Conservation in Turkmenistan" (2002) and "National Caspian Action Plan" (in preparation). No White-headed Duck have been ringed in Turkmenistan, although more than 50,000 other waterbirds have been marked as part of migration studies.

Table 15: Records of the White-headed Duck in Turkmenistan, 1980-2002.

Date	No. of birds recorded	Site name and location	Remarks	Counter/source
11 Feb 1980	3 in flight	Balkhan Bay, Caspian Sea		Vasilyev and Gauzer 2001
30 Nov 1980	930 in flight	Sarakamysh Lake	6 flocks	Vasilyev and Gauzer 2001
Nov 1981	450	Entire eastern Caspian coastline	21 small flocks	Vasilyev and Gauzer 2001
21-22 May 1982	2	Krasnovodsky Bay, southeast Caspian Sea	nesting site (1M+1F)	Vasilyev and Gauzer 2001
Jan 1983	1	Mainland lakes/rivers		Poslavski 1992
Dec 1984	64	Becovich Bay, southeast Caspian Sea	1 flock	Vasilyev and Gauzer 2001
Jan 1985	17	Mainland lakes/rivers		Poslavski 1992
Jan 1986	636	Mainland lakes/rivers		Poslavski 1992
Jan 1987	103	Mainland lakes/rivers		Poslavski 1992
1986-1988	100-200	Sarakamysh Lake	On migration	Poslavski 1992
1 Apr 1987	5	Sarakamysh Lake		Poslavski 1992
May 1987	5-6	Soltantagt Lake, Middle Amu Darya River	Broods of duckling	Poslavski 1992
Jan 1988	37	Krasnovodsky Bay, southeast Caspian Sea	1 flock	Vasilyev and Gauzer 2001
Jan 1988	358	Mainland lakes/rivers		Poslavski 1992
Summer 1989	19 pairs	Soltantagt Lake, Middle Amu Darya River		Poslavski 1992
Jan 1991	223	Krasnovodsky, Balhansky, Severo-Chelecensky Bays, southeast Caspian Sea	9 flocks	Vasilyev and Gauzer 2001
Jan 1993	3	Turkmenistan (unspecified)		Rose and Taylor 1993
14 Oct 1993	120 in flight	Tarta sea shoal, southeast Caspian Sea		Vasilyev and Gauzer 2001
28 Oct 1993	127 in flight	Balkhan Bay, Tarta Cape, southeast Caspian Sea		Vasilyev and Gauzer 2001
12-27 Oct 1994	313 in flight	Krasnovodsky Bay, southeast Caspian Sea	11 flocks (9-50 individuals)	Vasilyev and Gauzer 2001
18 Mar 1995	230 in flight	Krasnovodsky Bay, southeast Caspian Sea	6 flocks (14-43)	Vasilyev and Gauzer 2001
Jan 1998	820	Southeast coast of Caspian Sea, between Karabogasgol – Gasankuly		Vasilyev and Gauzer 2001
Jan 1998	171	Southeast coast of Caspian Sea, between Karabogasgol – Gasankuly		Vasilyev 2000
Jan 1999	7	Kianly Bay, southeast Caspian	1 flock	Vasilyev 2000
Nov 1999	268	Southeast coast of Caspian Sea, between Karabogasgol – Gasankuly		Vasilyev 2000
Jan 2000	19	Krasnovodsky Bay, southeast Caspian Sea	3 flocks (3,5,11)	Vasilyev 2001
Nov 2000	476	Southeast coast of Caspian Sea, between Karabogasgol – Gasankuly	9 flocks (9 –72)	Vasilyev 2001
Nov 2001	723	Southeast coast of Caspian Sea, between Karabogasgol – Gasankuly	16 flocks (7–55)	Vasilyev 2001
Jan 2002	21	Southeast coast of Caspian Sea, between Karabogasgol – Gasankuly	3 flocks (3–14)	V.I.Vasilyev, pers. comm., May 2002
Apr 2002	8	Krasnovodsky Bay, southeast Caspian Sea	At nesting site	M.E.Gauser, pers. comm., May 2002

Note: ¹ M - Male(s), F - Female(s)

Map 10: Distribution of the White-headed Duck in Turkmenistan during 1980-2002.



No.	Site Name	Location
1	Balkhan Bay, Caspian Sea	40°01'N, 52°48'E
2	Becovich Bay, southeast Caspian Sea	39°32'N, 52°50'E
3	Kianly bay, southeast Caspian Sea	40°12'N , 52°44'E
4	Krasnovodsky Bay, southeast Caspian Sea	39°42'-40°02'N, 52°53'-53°32'E
5	Sarakamysh Lake	41°50'-42°20'N, 56°52'-57°50'E
6	Southeast coast of Caspian Sea, between Karabogasgol - Gasankuly	37°23'-41°08'N, 52°60'-53°59'E
7	Severo-Chelecensky Bay, southeast Caspian Sea	39°33'-39°42'N, 53°14'-53°30'E
8	Soltantag Lake, southern Turkmenistan	38°47'N, 64°15'E

8.11.5 Threats

The threats reported to the White-headed Duck and its wetlands are based on information and feedback from government and non-government respondents to the survey.

- Hunting, poaching and inadequate capacity of conservation agencies.
- Increasing industrial and recreational pressure on wetlands, which have caused disturbance and degradation of the habitat.
- Drought in the last few years affecting water levels in wetlands.
- Lack of funds being allocated for research, awareness and conservation of waterbirds and wetlands.
- Lack of national experts in the field of species protection capable of providing quality information to manage wetlands and waterbirds.

8.11.6 Recommendations

The following recommendations have been proposed primarily on the basis of information and feedback from government and non-government respondents to the survey.

- Conduct regular surveys and monitoring programmes for the important habitats of wintering and breeding White-headed Duck.
- Establish protected areas for White-headed Duck and other migratory waterbirds at key sites along the coasts of the Caspian Sea and Amu Darya and Karakum Darya River Valleys.
- Enhance nature conservation and water management at national and local levels to ensure water availability for White-headed Duck and other migratory waterbirds.
- Produce and disseminate posters and other awareness materials related to the protection of the White-headed Duck and its key wetlands.
- Increase exchange of information on White-headed Duck and other waterbirds amongst researchers and government officials to enable improved conservation measures to be undertaken on the basis of up to date knowledge.
- Conduct training programmes for hunters, researchers, and managers and rangers of protected areas (nature reserves) on wetland and waterbird conservation, and identification of White-headed Duck and other waterbirds.
- Involve students in wetland and waterbird conservation activities to ensure greater grassroot support and awareness.

8.12 UZBEKISTAN

8.12.1 Historical records and distribution

In the past, the White-headed Duck was recorded as a breeding and passage migrant through Uzbekistan (Kashkarov 1987) (See Table 16). During northward migration, White-headed Duck were recorded in March on the Syr Darya River, near Chinaz Town and in April at Zeravshan and the Fergana Valley. During southward migration, they have been observed at the Khoresm Lakes, in the middle reaches of the Syr Darya River. There is only one historic record of wintering White-headed Duck in Uzbekistan – a bird found shot in the middle reaches of the Syr Darya River on 12 January 1966 (Kashkarov 1987). In recent years, small flocks of White-headed Duck have been found wintering in Aydarkul Lake (Nazarov unpublished).

A few historic records during the breeding season exist: one pair seen at a small saline lake near the northern foothills of the Nuratau mountains in July 1936 (Meklenburtcev 1937) and one bird was found shot at the Rogatoe Lake, near Uchkuduk Town in the Central Kyzylkum Desert on 17 August 1982 (Minaev 1987).

8.12.2 Recent records and distribution

Between autumn 1999 and summer 2001, comprehensive data on the ecology and numbers of White-headed Duck at the Sudochoye Wetlands have been collected by staff of the Institute of Zoology of Uzbek Academy of Sciences. Six expeditions have monitored numbers and distribution of White-headed Duck at the Sudochoye Wetlands during the recent drought in Central Asia.

Notable breeding and migrating populations of White-headed Duck have been discovered on Akushpa Lake, that forms a part of the Sudochoye Wetlands. A concentration of more than 3,000 individuals were also found during southward migration in 1999 (Kreuzberg-Mukhina and Lanovenko 2000, Kreuzberg-Mukhina *et al.* 2001). In mid April 2000, there were 1,166 birds at the lake and in July 2000, there were

more than 2,835 birds with 35 broods. However by autumn 2000, the effects of the drought (which began in the wetlands of Amu Darya Delta in 2000) started to affect White-headed Duck numbers. During autumn 2000, the number of White-headed Ducks was less than half of that in the previous year.

Table 16: Records of the White-headed Duck in Uzbekistan till 1980.

Date	No. of birds recorded	Site name and location	Counter/source
March 1888	Migrating flocks	Middle reaches of Syr Darya River, near Chinaz Town (40°54'N, 68°42'E)	Pleske 1888
	Rare	Middle reaches of of Amu Darya River, near Chardjou and Dargan-ata (40°28'N, 62°09'E)	Zarudny 1896
22-24 March 1903	Flocks	Zeravshan River, near Kermine and Ziatdin (40°08'N, 65°02'E)	Loudon 1910
4-10 April 1903	Migrating birds	Zeravshan River, near Kermine and Ziatdin	Loudon 1910
18 April 1903	Many	Fergana Valley, Station Melnikovo (40°10'N, 69°12'E)	Loudon 1910
July 1936	2 (one pair)	Saline lake on the northern foothills near Nuratau mountain (40°32'N, 67°25'E)	Meklenburctcev 1937
8-10 Oct 1957	Two flocks of 6 and 20 birds	Lower reaches of Amu Darya River, Khoresm district, lakes near Khazarasp Town (42°01'N, 61°01'E)	Salikhbaev and Bogdanov 1961
End Oct 1963	Single birds	Dalverzin Lakes, middle reaches of Syr Darya River (40°06' N, 69°04'E)	Kashkarov 1983
12 January 1966	1 bird shot	Dalverzin Lakes, middle reaches of Syr Darya River	Kashkarov 1987

Data collected from rangers, hunting inspectors and hunters, suggested the White-headed Duck left Akushpa Lake in November after it started to freeze over. The first White-headed Duck return to the lake at the end of February and beginning of March depending on when the ice thaws. During migration, White-headed Duck can be observed on other lakes of the Sudochoye Wetlands, but during the breeding season, all ducks only gather at the Akushpa Lake.

An overview of records of White-headed Duck in Uzbekistan is summarised in Table 17 and site locations are provided in Map 11.

8.12.3 Population and trends

At present, the population trend of White-headed Duck breeding in Uzbekistan in the southern section of the Aral Sea region depends mainly on climate conditions, especially the abundance of precipitation in Central Asia and management of the drainage system. Historically, local wetlands were supported by floodwaters from the Amu Darya River. The number of birds appears to depend on the natural conditions and management system of wetlands, but primarily on the irrigation system in the case of the wetlands in the Aral Sea region. Poor precipitation in the Aral region can not provide sufficient water for the stability of reservoirs (Evgeniya Lanovenko, pers. comm., May 2002; Elena Kreuzberg-Mukhina, pers. comm., May 2002).

The number of White-headed Duck had increased between 1999 and 2001, but have since declined. This decline is expected to continue for the next several years, because the main habitats are affected by natural and anthropogenic factors – drought and abstraction of water for agriculture. An increase in the number of White-headed Duck was noted in 1930 by N.A. Formozov (Isakov 1952), who observed that the number of White-headed ducks increased in the years of drought and decreased in the years with sufficient water which accounts for the distribution of the small groups within the small shallow lakes (Elena Kreuzberg-Mukhina, pers. comm., May 2002).

Table 17: Records of the White-headed Duck in Uzbekistan, 1980-2002.

Date	No. of birds recorded	Site name and location	Remarks	Counter/source
17 Aug 1982	1	Rogatoe Lake, near Uchkuduk Town, central Kyzylkum Desert	Bird shot	Minaev 1987
17 Oct 1999	40	Akushpa Lake, Sudochye Wetlands, southwest shore of Aral Sea	One flock, feeding in western part of the lake	Kreuzberg-Mukhina <i>et al.</i> 2001
20 Oct 1999	450	Akushpa Lake	Small flocks (7-30 birds)	Kreuzberg-Mukhina <i>et al.</i> 2001
21 Oct 1999	More than 3,000	Akushpa Lake	Small flocks (7-50) and two large flocks of 800+1500 birds	Kreuzberg-Mukhina <i>et al.</i> 2001
11 Jan 2000	1,137	Dengizkul Lake, S. Uzbekistan, near Turkmenistan border	Aerial count. Groups (575 birds, 200, 107, 175 and 85)	Lanovenko <i>et al.</i> 2000; Kreuzberg-Mukhina <i>et al.</i> 2001
3 Feb 2000	185	Dengizkul Lake	Aerial count. In one bay in small groups	Kreuzberg-Mukhina <i>et al.</i> 2001
5 Apr 2000	1 female	Aydar Lake	Last migratory species observed	Kreuzberg-Mukhina <i>et al.</i> 2001
8 Apr 2000	2 female	Tudakul Lake	Last migratory species observed	Kreuzberg-Mukhina <i>et al.</i> 2001
16-20 Apr 2000	1,166	Akushpa Lake	Small groups and pairs. Three flocks of up 90-230 birds	Kreuzberg-Mukhina <i>et al.</i> 2001
2-5 Jul 2000	2,835	Akushpa Lake	Females, ducklings, young non-flying individuals and flocks dominating males (500+800). Over 35 broods.	Kreuzberg-Mukhina <i>et al.</i> 2001
18-20 Oct 2000	1,370 birds	Akushpa Lake	Biggest flocks 230-380 birds.	Kreuzberg-Mukhina <i>et al.</i> 2001
2-5 April 2001	2,835 birds	Akushpa Lake	Small groups, about several hundred birds each. The largest flock of 600 birds.	Lanovenko <i>et al.</i> in preparation
3-6 July 2001	1,149	Akushpa Lake	In groups. Two flocks of 250-300. Broods were not recorded	Lanovenko <i>et al.</i> in preparation
19-23 Oct 2001	9	Karateren Lake, Akushpa Lake, Ustyurt collector, Sudochye Wetlands	Small groups. Karateren Lake 5 birds, Akushpa Lake 2, Ustyurt collector 2	Counted by Evgenia Lanovenko and Elena Kreuzberg-Mukhina, pers. comm. 2002
Nov 2001	Several tens	Aydar Lake	Small migratory flocks	Nazarov unpublished
26 Dec 2001	5	Tuzkan Lake and Aydar Lake	Feeding in the bays	Kreuzberg-Mukhina <i>et al.</i> 2002
27 Jan 2002	3	Dengizkul Lake	With other ducks	Kreuzberg-Mukhina <i>et al.</i> 2002
28 Jan 2002	6	Deukhona Lake	Flying flock	Kreuzberg-Mukhina <i>et al.</i> 2002
Nov - Feb 1997-2002	Several tens	Aydar Lake	Small flocks of 3-5 birds, 2 birds shot by hunters	Nazarov unpublished
7-17 Apr 2002	60	Akushpa Lake, Begdulla-Aydyn Lake, Ustyurt collector, Sudochye Wetlands	Migratory flocks. Akushpa Lake 27 birds, Begdulla-Aydyn Lake 30, Ustyurt collector 3.	Elena Kreuzberg-Mukhina, pers. comm. 2002
Oct 2002	700	Sudochye Wetlands		Elena Kreuzberg-Mukhina, pers. comm. 2002

Map 11: Distribution of the White-headed Duck in Uzbekistan during 1980-2002.



No.	Site name	Location
1	Sudochye Wetlands (Akushpa Lake, Karateren Lake, Begdulla-Ayдын Lake and Ustyurt collector)	58°30'N, 43°14'E
2	Aydar Lake	40°53'N, 66°35'E
3	Dengizkul Lake, south Uzbekistan, near border with Turkmenistan	39°07'N, 64°10'E
4	Deukhona lake	39°12'N, 64°39'E
5	Rogatoe Lake, near Uchkuduk town, Central Kyzylkum Desert	42°02'N, 63°37'E
6	Tudakul Lake	39°52'N, 64°46'E
7	Tuzkan Lake	40°37'N, 67°11'E

8.12.4 Conservation status

The White-headed Duck is protected under the Law of the Republic of Uzbekistan on protection and usage of animals (December 1997).

The White-headed Duck is included in the national Red Data Book (1983) as a “probably extinct species” in Central Asia. With the recent observations on the status of the species, in the new edition of the Red Data Book (in press), it is to be listed as an Endangered species. The Red Data Book of Uzbekistan is a judicial document, which provides direct protection for all listed species.

The national hunting regulations prohibit the hunting of White-headed Duck, with high fines being imposed for the violation of this regulation.

The most important sites for White-headed Duck in Uzbekistan are the Sudochye Wetlands and Dengizkul Lake. Both these sites are non-hunting areas or "Zakaznik" (this designation corresponds to IUCN category IV for protected areas). Additionally, Dengizkul Lake was designated as a Ramsar Site in February 2002.

The process of producing regional and national action plans on the threatened species is now ongoing, and it is proposed that based on a regional action plan, a national action plan will be produced.

8.12.5 Threats

Recent work in Sudochoye Wetlands and other sites by Elena Kreuzberg-Mukhina, Evgenia Lanovenko and colleagues have provided up to date information on current threats to the White-headed Duck in Uzbekistan. Their findings are summarised below:

- **Changes in hydrological regime:** The key sites for White-headed Duck in Uzbekistan have no stable hydrological regime. The Sudochoye Wetland is under threat of drying out completely due to a combination of the change in the water-regime in the Aral Sea basin and the ongoing extended drought in Central Asia. In the summer of 2001, the Sudochoye Wetland was completely dry while Akushpa Lake had changed into a swamp. In the winter and spring of 2002, the lake received some water as precipitation but drainage water arrived into the lake only in July. So, in the breeding season of 2002 there was not enough water in the lake and young reeds did not grow. As a result of this, the lake could not provide sufficient habitats for the White-headed Duck. Historically, during the 20th century, the Sudochoye Wetland has some times dried up and it has changed from a fresh water lake to contain saline water. Dengizkul Lake has turned saline and is drying up. Changes have taken place in the drainage system which is feeding water into this lake. These changes are linked with the implementation of a large-scale project, the aim of which is to improve the drainage system in Uzbekistan and water quality in Amu Darya River (and for irrigation, as outlined below).
- **Over-abstraction of water:** The Sudochoye Wetland is under threat of drying out completely due to a combination of the change in the water-regime in the Aral Sea basin and the ongoing extended drought in Central Asia. In the winter of 2001, the Sudochoye Wetland was completely dry and Akushpa Lake had changed into a swamp. Dengizkul Lake has turned saline and is drying up, because the water sources that fed it are being used for irrigation purposes in the desert villages located close to the Amu Darya River.
- **Climatic effects:** It is possible that future natural climatic cycles, exacerbated by the effects of global warming, will have an impact the ecological conditions of the White-headed Duck habitat.
- **Burning of reed beds:** Burning of reed beds in the spring season is also a major threat to the habitat of White-headed Duck and other reed-nesting species. The main reason for reed burning by local people is to clean up old plants and to ensure the better growth of young ones, which they use as cattle fodder.

8.12.6 Recommendations

The following recommendations have been proposed primarily on the basis of information and feedback from government and non-government respondents to the survey.

- Develop a national action plan for the White-headed Duck.
- Conduct surveys at potential sites to determine a more accurate status of the White-headed Duck in Uzbekistan.
- Promote wise use of water resources at the two most important wetlands for White-headed Duck.
- Create specially protected areas at key breeding, migratory and staging sites for the White-headed Duck in Uzbekistan.
- Improve the monitoring and management system of wetlands in Uzbekistan to ensure that qualitative and quantitative information can be obtained on an ongoing basis.
- Distribute information on the conservation of White-headed Duck through the media and through local education campaigns.
- Undertake awareness raising initiatives amongst local hunters to increase their understanding of the threatened status of White-headed Duck and other species and the need for their conservation.

- Undertake an education programme for rangers and inspection staff on wetlands and waterbirds to raise awareness and skills to conserve threatened species.
- Increase capacity among nature protection agencies and other conservation bodies (NGOs, local communities, etc.) to ensure conservation work on wetlands and waterbirds can be undertaken at the national and local levels.

REFERENCES

- *Abdurakhmanov, G. M. 1998. *Red Data Book of the Republic of Daghestan*. Makhachkala. (in Russian)
- *Abdusalamov, I. A. 1971. *Birds of Tajikistan* (Fauna of Tadjik SSR, volume 19, part 1). Institute of Zoology and Parasitology of Academy of Sciences of Tadjik SSR, Dushanbe. (in Russian)
- *Ali, S. 1936. Economic Ornithology in India. *Curr. Sci.* 4: 472–478.
- Ali, S. and Ripley, S. D. 1983. *Handbook of the Birds of India and Pakistan*. Oxford University Press, New Delhi.
- Al-robaae, K. H. and Salem, Y. A. 1996. In *Gibier Faune Sauvage, Game Wildlife 13* (ed. Birkan, M., van Vessem, J., Havet, P., Madsen, J., Trolliet, B. and Moser, M.), Proceeding of the Anatidae 2000 Conference, Strasbourg, France, 5-9 December 1994. pp 275-283. Paris.
- *Amat, J. A. and Sanchez, A. 1982. Biología y ecología de la Malvasia *Oxyura leucocephala* Andalucía. *Donana Acta Vertebr.* 9: 251-320. (in Spanish)
- *Anon. 1985. *Red Data Book of Kirghiz SSR*. “Kyrgyzstan” Publishers: Frunze. (in Russian)
- *Anon. 1988. *Red Data Book of Tajikistan*. Donish, Dushanbe. (in Russian)
- Anon. 1997a. News of Israel. *Threatened Waterfowl Specialist Group News* 10: 6.
- Anon. 1997b. News of Turkey. *Threatened Waterfowl Specialist Group News* 10: 7.
- Anstey, S. 1989. *The Status and Conservation of the White-headed Duck Oxyura leucocephala*. IWRB Special Publication No. 10. International Waterfowl and Wetlands Research Bureau, Slimbridge.
- *Ataev, C. A., Vasiliyev, V. I. and Gorelova, R. N. 1978. *Studies of infrequent and endangered species of birds of Turkmenistan*. Academy of Sciences of Turkmenistan, Series of Biological Sciences 4. Ashkhabad. (in Russian)
- Azafzaf, H. 2001. White-headed Duck in Tunisia. *Threatened Waterfowl Specialist Group News* 13: 37.
- *Baranov, A. A. 1991. *Rare and insufficiently known birds in Tuva*. Krasnoyarsk University Press, Krasnoyarsk. (in Russian)
- Behrouzi-Rad, B. 1996. Threatened Anatidae Species in Iran. In *Gibier Faune Sauvage, Game Wildlife 13* (ed. Birkan, M., van Vessem, J., Havet, P., Madsen, J., Trolliet, B. and Moser, M.), Proceeding of the Anatidae 2000 Conference, Strasbourg, France, 5-9 December 1994. pp 815-829. Paris.
- *Belyalov, O. V., Annenkova, S. Y. and Karpov, F. F. 2002. *New data about White-headed Duck on the south and Southeast of Kazakhstan*. Zoological Researches in Kazakhstan, Almaty. (in Russian)
- *Berezovikov, N. N., Gubin, B. M., Gul, I. R., Erokhov, S. N., Karpov, F. F. and Kovalenko, A. V. 1999. *Birds of the Taukumy desert (southeastern Kazakhstan)*. Academy of Medicine and Biological Sciences, Publ. House of Ukrainian, Kiev. (in Russian)
- BirdLife International. 2000. *Threatened Birds of the World*. Lynx Edicions and BirdLife International, Barcelona and Cambridge.
- BirdLife International. 2001. *Threatened Birds of Asia: the BirdLife International Red Data Book*. BirdLife International, Cambridge.

- *Boldbaatar Sh. 1997. White-headed Duck. In *Mongolian Red Book* (ed. Shiirevdamba, T.), pp 103-104. Ministry for Nature and the Environment of Mongolia, Ulaanbaatar. (in Mongolian)
- *Braude, M. I. 1989. *The modern fauna of Anseriformes of the Middle Ural. Distribution and fauna of birds of the Ural*. Informative Materials. Uralian Branch of Academy of Science of the USSR, Sverdlovsk. (in Russian)
- Bräunlich, A. and Tseveenmyadag, N. *Status and conservation of the White-headed Duck Oxyura leucocephala in Mongolia*. (in preparation)
- Bräunlich, A. 1995. *Report on the first WWF-Expedition to the Great Lakes Basin, Western Mongolia, May - July 1995, and preliminary recommendations for the establishment of a new protected area*. WWF-Mongolia, Ulaanbaatar. (unpublished)
- Buckley, Y., Holt, L., Pullen, T., Robinson, K., Boyla K. and Can, O. 1998. Breeding Status of the White-headed Duck on the Central Plateau, Turkey. *Threatened Waterfowl Specialist Group News* 11: 35-36.
- *Bukreev, S. A. and Chernobay, V. F. 2001. Current status of the White-headed Duck in the Lower Volga region. In *Problems of study and protection of the Anseriformes birds of Eastern Europe and Northern Asia*. pp 25-26. Moscow. (in Russian)
- *Bukreev, S. A. and Chernobay, V. F. 2002. Monitoring of European Russia IBAs: Sarpinskiye Lakes. *Russian IBA Newsletter* 15: 7. (in Russian)
- Chaudhry, A. A. 2002. *White-headed Duck Survey in Pakistan: 2002*. Wetlands International, Kuala Lumpur. (unpublished)
- *Cheng Tso-hsin. 1987. *A synopsis of the avifauna of China*. Science Press, Beijing.
- Cresswell, W., Yerokhov, S., Berezovikov, N., Mellanby, R., Bright, S., Catry, P., Freile, J., Gretton, A., Zysin, A., McGregor, R. and McLaughlin, D. 1999. Important Wetlands in Northern and Eastern Kazakstan. *Wildfowl* 50: 181-194.
- Criado, J. 1997. Urgent Action Needed to save the White-headed Duck. *Threatened Waterfowl Specialist Group News* 10: 16-17.
- del Hoyo, J., Elliott, A. and Sargatal, J. eds (1992). *Handbook of the Birds of the World*. Vol. 1. Lynx Edicions, Barcelona.
- Delany, S., Reyes, C., Hubert, E., Pihl E., Haanstra, L. and Van Strien, A. 1999. *Results from the International Waterbird Census in the Western Palearctic and Southwest Asia, 1995 and 1996*. Wetlands International Publication No. 54, Wageningen.
- *Dementiev, G. P. and Gladkov, N. A. 1952. *Birds of the Soviet Union*, Vol. 4. 1967 translation, Israel Program for Scientific Translation, Jerusalem.
- Dimitrov, M., Profirov, L., Nyagolov, K. and Michev, T. 2000. Record Counts of White-headed Duck in Bulgaria. *Threatened Waterfowl Specialist Group News* 12: 18-20.
- *Dolgushin, I. A. 1960. *Ptisty Kazakhstana* 1. Alma-Ata. (in Russian)
- *Dzhamirzoev, G. S. 2002. Monitoring of European Russia IBAs: Adzhi Lake. *Russian IBA Newsletter* 15: 14-15. (in Russian)
- Eken, G. 1998a. Status of the Globally Threatened Wildfowl on Central Anatolian Lakes, Turkey. *Threatened Waterfowl Specialist Group News* 11: 32-33.

- Eken, G. 1998b. Status of the Globally Threatened Wildfowl at Burdur Golu, Turkey. *Threatened Waterfowl Specialist Group News* 11: 10.
- Evans, M. I. 1994. *Important Bird Areas in the Middle East*. BirdLife International, Cambridge.
- Gilissen, N., Haanstra, L., Delany, S., Boere, G. and Hagemeyer, W. 2002. *Numbers and distribution of wintering waterbirds in the Western Palearctic and Southwest Asia in 1997, 1998 and 1999. Results from the International Waterbird Census*. Wetlands International Global Series No. 11, Wageningen.
- *Gordienko, N. S., Drobvtsev, V. I. and Koshelev, A. I. 1986. Biology of White-headed Duck in Northern Kazakhstan and south of West Siberia. In *Rare, threatened and little known birds of USSR Collection articles*, pp 8-15. Central Board for Nature Conservation of the RSFSR, Central Science Research Laboratory, Moscow. (in Russian)
- *Gordienko, N. S. 1995. On distribution of rare species of waterfowl of Southern Trans-Uralia. In *Materials on distribution of birds in the Ural, Cis-Uralia and West Siberia: informative Materials* (ed. Ryabitsev, V. K.), pp 15-16. Uralian Branch of Russian Academy of Science, Moscow. (in Russian)
- *Gordienko, N. S. 1998. The modern status and tactics of conservation of rare bird species of Southern Trans-Uralia. In *Rare bird species of non-black earth center of Russia*, pp 294-298. Moscow. (in Russian)
- *Gordienko, N.S. 2001. *Waterbirds in Southern Ural*. Akademii of Ural department, Miass. (in Russian)
- Green, A. J. 1992. *The Status and Conservation of the White-winged Wood Duck Cairina scutulata*. IWRB Special Publication No. 17, IWRB, Slimbridge.
- Green, A. J. and Anstey S. 1992. The status of the White-headed Duck *Oxyura leucocephala*. *Bird Conservation International* 2: 185-200.
- Green, A. J. and Hughes, B. 1996. Action plan for the White-headed Duck *Oxyura leucocephala*. In *Globally threatened birds in Europe* (ed. Heredia, B. Rose, L. and Painter, M.), pp 119-146. Council of Europe Publishing, Strasbourg.
- Green, A. J. and Hughes, B. 2001. White-headed Duck *Oxyura leucocephala*. In *BWP Update: the journal of birds of the Western Palearctic*, Vol. 3, No. 2 (ed. D.B. Parkin), pp 79-90. Oxford University Press, Oxford.
- Green, A. J. and Hunter, J. 1996. The Declining White-headed Duck: A call for information. *Threatened Waterfowl Specialist Group News* 9: 19.
- Green, A. J. and Yazar, M. 1996. Rapid decline of White-headed Duck at Burdur Lake, Turkey. *Threatened Waterfowl Specialist Group News* 9: 16-18
- *Guan Guanxun and Cheng Tso-hsin. 1962. *Oxyura*, a new genus of bird to China. *Acta Zool. Sinica* 14(3). (in Chinese)
- Handrinos, G. I. 1995. White-headed Duck in Greece. *Threatened Waterfowl Specialist Group News* 7: 6-7.
- Handrinos, G. I. 1998. Record Count of White-headed Duck in Greece. *Threatened Waterfowl Specialist Group News* 11: 34-35.
- Heinicke, T. 2001. White-headed Ducks in Central Kazakhstan. *Threatened Waterfowl Specialist Group News* 13: 7.

- Hughes, B. 1992. *The ecology and behaviour of the Ruddy Duck Oxyura jamaicensis jamaicensis Gmelin in Great Britain*. Ph. D. Thesis, University of Bristol.
- Hughes, B. 1996. The Ruddy Duck (*Oxyura jamaicensis*) in Europe and the Threat to the White-headed Duck (*Oxyura leucocephala*): A Review, an Evaluation and Conservation Actions. In *Gibier Faune Sauvage, Game Wildlife 13* (ed. Birkan, M., van Vessem, J., Havet, P., Madsen, J., Trolliet, B. and Moser, M.), Proceeding of the Anatidae 2000 Conference, Strasbourg, France, 5-9 December 1994. pp 1127-1141. Paris.
- *Hume, A. O. 1887. Catalogue of the birds in the Provincial Museum, N. W. P and Oudh, Lucknow. *Stray Feathers* 10: 442-444.
- International Research Institute for Climate Prediction. 2001. *The Drought and Humanitarian Crisis in Central and Southwest Asia: A Climate Perspective*. IRI Special Report 01-11. New York.
- *Irisov, E. A. and Irisova, N. L. 1982. Records of some birds, listed in Red Data book of the USSR in the territory of Altai Krai. In *Endangered and rare plants and animals of the Altai Territory and problems of their conservation: abstracts of Conference*. pp 45-47. Barnaul. (in Russian)
- *Isakov, Y. A. and Vorobyev C. A. 1940. *The browse of winterings and flyover of birds on an Austral Caspian*. Transactions of the All-Union Ornithological Reservation GasanKuly, discharge 1, Moscow.
- *Isakov, Y. A. 1952. Order Anseriformes. Family Anatidae. *Birds of Soviet Union* (ed. Dementiev, G. P. and Gladkov, N.A.). pp 344-636. Publ. House "Soviet Science", Moscow.
- Islam, M. Z. and Rahmani, A. 2001. *White-headed Duck (Oxyura leucocephala) in Amakhera Wetlands, Aligarh District, Uttar Pradesh, India*. Bombay Natural History Society, Mumbai. (unpublished)
- IUCN. 2000. *IUCN Red List of Threatened Species*. IUCN-The World Conservation Union, Gland.
- *Ivanov, A. I. 1940. *Erismatura leucocephala* (Scop.). In *Birds of Tajikistan*. pp 46. Proceedings of Tajik Base of the Academy of Sciences of the USSR. Vol. 10, Zoology and Parasitology. Publishing House of the Academy of Sciences of the USSR, Moscow. (in Russian)
- *Ivanov, A. I. 1969. White-headed Duck – *Oxyura leucocephala* (Scop.). In *Birds of Pamiro-Alai*. pp 57. Zoological Institute of the Academy of Sciences of the USSR. "Nauka", Leningrad. (in Russian)
- *Jamirzoev, G. S. 1999. *Territorial Protection of migratory birds in Daghestan. Inventarization, monitoring and protection of the Important Bird Area of Russia*. Russian Bird Conservation Union, Moscow. (in Russian)
- *Johansen, H. 1959. Die Vogelfauna Westsibiriens. III. Teil (Non-Passeres), 7. Fortsetzung: Anseres II (*Anas-Mergus*). *J. Ornithol.* 100: 313-36.
- *Karyakin, I. V. and Kozlov, A. A. 1999. *Preliminary cadastre of birds of Chelyabinsk Region*. "Manuscript" Press, Novosibirsk. (in Russian)
- *Karyakin, I. V. 1998. *Conspectus of bird fauna of Republic of Bashkortostan*. Center of Field Investigations of Uralian Animal Conservation Union, Perm. (in Russian)
- *Karyakin, I. V., Bystrykh, S. V. and Konovalov, L. I. 1999. *Ornithofauna of the Sverdlovsk Region*. Novosibirsk. (in Russian)
- *Kashkarov, D. Y. 1983. The White-headed Duck. In *Red Data Book of the Uzbek SSR*. Vol.1. Vertebrate Animals. pp 66-68. Fan. Tashkent. (in Russian)

- *Kashkarov, D. Y. 1987. The White-headed Duck. In *Birds of Uzbekistan*. Vol. 1. pp.115-116. Fan. Tashkent. (in Russian)
- *Korshikov, L. V. and Kornev, S. V. 1999. News of the 1999 ornithological season in Orenburg Territory. In *Materials on bird distribution in the Urals, in Cis-Ural area and West Siberia*. pp.140-142. Ekaterinburg. (in Russian)
- *Kovshar (ed.) 1996. *Red Data Book of Kazakhstan*. Vol.1, Animals, Part 1. Vertebrates. "Konzhik", Almaty. (in Russian)
- *Kovshar, A. F. and Berezovikov, N. N. 2000. *Ornithological observations in Naurzum (Northern Kazakhstan) in spring 1998 and 1999*. In (Materials on distribution of birds in Ural, in Cis-Uralia and west Siberia). Uralian Branch of Russian Academy of Sciences, Ekaterinburg. Pp 94-114. (in Russian)
- Kreuzberg-Mukhina, E., Kashkarov, D., Lanovenko, E., Nazarov, E., and Shernazarov, E. 2000. Status of threatened Anatidae in Uzbekistan. *Threatened Waterfowl Specialist Group News* 12: 70-75.
- Kreuzberg-Mukhina, E. and Lanovenko, E. 2000. White-headed Duck at the Sudochoye Wetlands, Uzbekistan. *Threatened Waterfowl Specialist Group News* 12: 15-16.
- Kreuzberg-Mukhina, E., Lanovenko, E., Filatov, A., and Zagrebin, S. 2001. Status and Distribution of the White-headed Duck in Uzbekistan. *Threatened Waterfowl Specialist Group News* 13: 46-48.
- Kreuzberg-Mukhina, E., Lanovenko, E. and Shernazarov, E. 2002. *Status survey and preliminary assessment of migratory and wintering places of the Lesser White-fronted Goose in Uzbekistan*. (unpublished)
- Krivenko, V.G., (ed.) 1999. *Wetlands in Russia, Volume 1: Wetlands of International importance*. Wetlands International, Moscow.
- Krivenko, V.G., (ed.) 2000. *Wetlands in Russia, Volume 3: Wetlands on the Ramsar Shadow List*. Wetlands International, Moscow.
- Kirwan, G. M. 1995. Breeding Status of the White-headed Duck in Turkey. *Threatened Waterfowl Specialist Group News* 8: 14-15.
- *Kuchin, A. P. 1991. *Rare animals of Altai*. NGPI, Novosibirsk. (in Russian)
- Lachmann, L., Eichhorn, G., Heinicke, T., Ratayczak, J. and Schielzeth, H. 2002. *Positive findings about White-headed Duck population in the Tengiz Region/Central Kazakhstan*. (in preparation)
- *Ladhar, S. S., Chauhan, M., Handa, S. M. and Jerath, N. 1994. *Harike Lake, Punjab*. WWF-India, Delhi.
- Lanovenko E., Filatov A. and Zagrebin S. 2000. White-headed Duck at Dengizkul Lake, Uzbekistan. *Threatened Waterfowl Specialist Group News* 12: 16.
- *Laptev, M. K., Sulima, V. I. and Freyberg, L. P. 1934. *All-Union Ornithological reservation in GasanKuly Turkmenish SSR*. The report of Turkmenish Interdepartmental Committee on Nature Protection and Development of Natural Resources, Aschcabat. (in Russian)
- Liegl, C. 1998. *Waterfowl at Har Us Nuur National Park and at Ayrag Nuur (Western Mongolia) - Report on two Expeditions in June and September 1998*. WWF-Mongolia, Ulaanbaatar. (unpublished)
- *Linkov, A. B. 1984. On the White-headed Duck ecology in Eastern Manych. In *Modern status of waterfowl resources*. pp 85-86. Moscow.

- *Linkov, A. B. 1984. On ecology of the White-headed Duck on the Eastern Manych. In *Modern status of waterfowl resources* (Abstracts of All-Union Meeting 20-23 Oct 1984). pp 85-86. Moscow.
- *Linkov, A. B. 2001. White-headed Duck. In *Red Data Book of the Russian Federation (Animals)*. pp 418-419. ACT Press and Astrel' Press, Moscow. (in Russian)
- Lopez, A and Mundkur, T. eds. 1997. *The Asian Waterfowl Census 1994-1996. Results of the Coordinated Waterbird Census and an Overview of the Status of Wetlands in Asia*. Wetlands International, Kuala Lumpur.
- *Loudon, H. 1909, 1910. Meine dritte Reise nach Zentral-Asien und ihre ornithologische Ausbeute. – *J. Ornith.*, Jg. 57: .505-573; Jg. 58: 1-90. (in German)
- Ma Ming 2001. *A Checklist of the Birds in Xinjiang, China*. Arid Zone Research, 18 (Supplement). Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences. Urumqi. (in Chinese)
- MacKinnon, J., Phillipps, K., and He, F. Q. 2000. *A Field Guide to the Birds of China*. Oxford University Press. Changsha. (in Chinese)
- *McCracken, K. G., Harshman, J., McClennan, D.A. and Afton, A.D. 1999. Data set incongruence and correlated character evolution: an example of functional convergence in the hind-limbs of stiff-tail diving ducks. *Syst. Biol.* 48(4): 683-714.
- *Meklenburtsev, R. N. 1937. *Materials on the fauna of birds and mammals of the Nuratau range*. Transactions of Samarkand University, V. VIIIa, issue 26. (in Russian)
- *Minaev, N.A. 1987. Some bird species in Central Kyzylkum desert. In *Mammals and birds of Uzbekistan*, pp 92-93. Fan, Tashkent. (in Russian)
- Morozov, V. and Shevchenko, V. 1998. White-headed Duck in the North Caspian Region. *Threatened Waterfowl Specialist Group News* 11: 36.
- *Moseikin, V. N. 2000. New ornithological findings in Saratov Region. *Russ. J. of Ornithology* 104: 3-7.
- Mundkur, T. and Taylor, V. 1993. *Asian Waterfowl Census 1993*. AWB, Kuala Lumpur and IWRB, Slimbridge.
- Munteanu, D. 2000. White-headed Duck in Romania. *Threatened Waterfowl Specialist Group News* 12: 17.
- Munteanu, D. 1995. Current Status of the White-headed Duck in Romania. *Threatened Waterfowl Specialist Group News* 7: 7-8.
- Panayotopoulou, M. and Green, A. J. 2000. White-headed Duck in Greece. *Threatened Waterfowl Specialist Group News* 12: 16.
- *Pandit, A. K. 1982. Looking out for the wildlife in Jammu and Kashmir Himalayas. *Environmental Awareness* 5: 52–61.
- Perennou, C., Rose, P. and Poole, C. 1990. *Asian Waterfowl Census 1990*. IWRB, Slimbridge.
- Perennou, C. and Mundkur, T. 1991. *Asian Waterfowl Census 1991*. IWRB, Slimbridge.
- Perennou, C. and Mundkur, T. 1992. *Asian Waterfowl Census 1992*. IWRB, Slimbridge.

- Perennou, C., Mundkur, T., Scott, D. A., Follestad, A. and Kvenild, L. 1994. *The Asian Waterfowl Census 1987-91: Distribution and Status of Asian Waterfowl*. AWB Publication No. 86. Asian Wetland Bureau, Kuala Lumpur.
- *Pleske, T. 1888. *Revision der turkestanischen Ornithologie: Nach Sammlungen des verstorbenen Konservators V. Russow*. St.-Pb., p58 (Mem. Acad. sci. St.-Pb. Ser.7, Vol. 36, N3). (in German)
- Poslavski, A. 1992. Status of Marbled Teal and White-headed Duck in Turkmenistan and Uzbekistan. *Threatened Waterfowl Specialist Group News* 2: 8-10.
- Profirov, L. and Dimitrov, M. 2001. White-headed Duck at Lake Vaya, Bulgaria. *Threatened Waterfowl Specialist Group News* 13: 9.
- Rogacheva, E. V. 1992. *The Birds of Central Siberia*. Husum Druck.
- Roberts, T. J. 1991. *Birds of Pakistan*. V1. Oxford University Press. Karachi.
- Rose, P. M. 1992. *Western Palearctic Waterfowl Census 1992*. IWRB, Slimbridge.
- Rose, P. M. (Ed.). 1995. *Western Palearctic and South West Asia Waterfowl Census 1994*. IWRB, Slimbridge.
- Rose, P. M. and Scott, D. A. 1997. *Waterfowl Population Estimates - Second Edition*. Wetlands International Publ. 44, Wageningen.
- Rose, P. M. and Taylor, V. 1993. *Western Palearctic and South West Asia Waterfowl Census 1993*. IWRB, Slimbridge.
- *Salikhbaev, K. S. and Bogdanov, A. N. 1961. *Birds*. Part 3. Publishing House of Academy of Science of the UzSSR. Tashkent. (Fauna of Uzbek SSR / Ac of Sc of UzSSR. Inst. of Zool. and Parasitology; V.2). (in Russian)
- *Sanchez, M. I., Green, A. J., and Dolz, C. 2000. The diets of the White-headed Duck *Oxyura leucocephala*, Ruddy Duck *O. jamaicensis* and their hybrids from Spain. *Bird Study* 47: 275-284.
- *Savage, C. D. W. 1965. White-headed Ducks in Pakistan. *Wildfowl Trust Ann. Rep.* 16: 121-126.
- Scott, D. A. (ed.) 1989. *A Directory of Asian Wetlands*. IUCN, Gland and IWRB, Cambridge.
- Scott, D. A. (ed.) 1995. *A Directory of Wetlands in the Middle East*. IUCN, Gland and IWRB, Slimbridge.
- Scott, D. A. and Rose, P. M. 1989. *Asian Waterfowl Census 1989*. IWRB, Slimbridge.
- Scott, D. A. and Rose, P.M. 1996. *Atlas of Anatidae Populations in Africa and Western Eurasia*. Wetlands International Publication No.41, Wetlands International, Wageningen.
- Sebastian, A. and Davies, J. 2001. *East Dongting Lake National Nature Reserve, Hunan Province - Biological Resource Assessment, Annotated Checklist of Vertebrate Species*. Report to UNDP/GEF Project Wetland Biodiversity Conservation and Sustainable Use in China. Beijing. (unpublished)
- Shahbazi, Y. 2000. Status of the White-headed Duck in Eastern Azarbijan, Iran. *Threatened Waterfowl Specialist Group News* 12: 34-38.
- *Shubin, A. O. 2001. Monitoring of European Russia IBAs: Sarpinskiye Lakes. *Russian IBA Newsletter* 13: 18. (in Russian)
- *Singh, G. 1992. Status of Harike wetland. *Tigerpaper* 19(3): 26-28.

- Sutanov, E. H. 2001, Status of the White-headed Duck in Azerbaijan. *Threatened Waterfowl Specialist Group News* 13: 44-45.
- Syroechkovski E. E. and E. V. Rogacheva. 1995. Red Data Book of the Krasnoyarsk Territory. Rare and Endangered Species of Animals. Krasnoyarsk Book Publishing House.
- Torres, J. 2001. White-headed Duck in Spain in 2000. *Threatened Waterfowl Specialist Group News* 13: 7.
- Torres, J. 2001. New Records of White-headed Duck from Morocco. *Threatened Waterfowl Specialist Group News* 13: 43.
- Torres, J. 2000. White-headed Duck in Spain. *Threatened Waterfowl Specialist Group News* 12: 15.
- *Toress, J. A. and Moreno-Arroyo, B. 2000a. La recuperacio'n de la malvasi'a cabeciblanca (*Oxyura leucocephala*) en Espan~a durante el u'ltimo decenio del siglo XX. *Oxyura* 10 (in Spanish)
- *Toress, J. A. and Moreno-Arroyo, B. 2000b. Presencia de la Malvasi'a canela (*Oxyura jamaicensis*) en Espan~a. *Oxyura* 10. (in Spanish)
- *Urdiales, C. and Pereira, P. 1993. *Identification key of O. jamaicensis, O. leucocephala and their hybrids*. ICONA, Madrid.
- Van der Ven, J. 1987. *Asian Waterfowl 1987. Midwinter bird observations in some Asian countries*. IWRB, Slimbridge.
- Van der Ven, J. 1988. *Asian Waterfowl 1988. Midwinter bird observations in most Asian countries*. IWRB, Slimbridge.
- Van der Ven, J. 2002. *Looking at birds in Kyrgyz Republic*. Rarity Publishers, Bishkek.
- *Vasiliyev, V. I., Caravaev, A. A. and Rustamov E. A. 1984. *Allocation and number of birds on the winterings Turkmenish S.S.R., The report " Ecologo - faunistic aspects of analysis of the fauna of Turkmenistan"*. Ashcabat.
- Vasil'yev, V. I. 1999. White-headed Duck. In *Red Data Book of Turkmenistan*, Vol 1, pp 222-223. Turkmenistan Press, Ashgabat. (in Turkmenian and in Russian).
- *Vasilyev, V. I. 2000. In "The chronicle of the Nature Hasar State reserve for 2000", volume 26, Turkmenbashy situ. (unpublished).
- *Vasilyev, V. I. 2001. In "The chronicle of the Nature Hasar State reserve for 2001", volume 27, Turkmenbashy situ. (unpublished).
- Vasilyev, V. I. and Gauser, M. 2001. Status of threatened Waterfowl in the Southeast Caspian Region of Turkmenistan. *Threatened Waterfowl Specialist Group News* 13: 69-71
- Vasilyev, V. I. and Gauser, M. 2001. Status of Waterbirds in the Caspian Region of Turkmenistan, 1970-2000. *Threatened Waterfowl Specialist Group News* 13: 72-73
- Wang Sung, Zheng Guangmei and Wang Qishan. 1998. *China Red Data Book of Endangered Animals-Aves*. Science Press, Beijing. (in Chinese)
- Wetlands International. 2002. *Waterbird Population Estimates – Third Edition*. Wetlands International, Global Series No. 12, Wageningen.

- *Whitehead, W. A. 1931. Notes on the White-headed Duck or Stiff-tail (*Erismatura leucocephala*). *Journal of the Bombay Natural History Society* 35(1): 211-212.
- *Yakimenko, V. V. 1998. Materials on bird distribution in Omsk Region. *Materials on bird distribution in the Urals, in Cis-Ural area and West Siberia*. pp 192-221. Ekaterinburg. (in Russian)
- *Zabelin, V. I., Munhtogtokh, O. and Bayarkhuu, C. 2000. To the current status and perspectives for protection of rare birds in Tuva and Northwest Mongolia. In *Inventory, monitoring and conservation of important bird areas in Russia*. pp. 100-104. Moscow. (in Russian)
- *Zarudny, N. A. 1896. Ornithological fauna of the Transcaspian Territory (Northern Persia, Transcaspian region, Khiva oasis and the plain Bukhara). In *Materials on study of fauna and flora of the Russian Empire*. Issue II. Pp 1-555. Dept. Zool., Moscow.
- * Zoological Survey of India. 1994. *Red Data Book on Indian Animals: Part 1: Vertebrata* (Mammalia, Aves, Reptilia and Amphibia). Zoological Survey of India, Calcutta.
- *- references not seen in original

ADDITIONAL READING

- Anstey, S. and Moser, M. E. 1990. A recovery plan for the White-headed Duck. In *Managing waterfowl populations* (ed. G.V.T. Matthews). IWRB Special Publication 12. pp 204-206. Slimbridge.
- Asia-Pacific Migratory Waterbird Conservation Committee. 2001. *Asia-Pacific Migratory Waterbird Conservation Strategy: 2001-2005*. Wetlands International - Asia Pacific, Kuala Lumpur.
- Atadjanov, A., Filatov, A., Lanovenko, Y., Zagrebin, S. Chernogaev, E. and Khodijaev, J. 2001. *Aerial Survey of Wetlands in Uzbekistan (Winter 2000)*. Part 3 of Protection of Uzbekistan's Wetlands and their Waterfowl. Tashkent. (unpublished).
- Baker, E. C. S. 1908. *Indian ducks and their allies*. Bombay Natural History Society, Mumbai.
- Borodin, A. M. 1984. *Red Data Book of the USSR*. Moscow.
- Bukreev, S. A. 1997. *Ornithogeography and Nature Protection in Turkmenistan*. Moscow. (in Russian)
- Cunningham, A. H. 1928. Notes on duck shooting in the Roorkee district, U.P., in the years 1903 to 1927. *J. Bombay Nat. Hist. Soc.* 32: 600–605.
- Chaudhry, A. A., Gill, A. H. and Ali, Z. 1997. *Conservation of White-headed Duck in the Salt Range Lakes (Ucchali Complex), Punjab-Pakistan*. Punjab Wildlife Research Institute, Faisalabad.
- Chaudhry, A. A. 1992. Habitat Changes Threaten the White-headed Duck in Punjab, Pakistan. *Threatened Waterfowl Specialist Group News* 2: 5-6.
- Green, A. J., Hilton, G., Hughes, B., Fox, A. D. and Yarar, M. 1993. The White-headed Duck at Burdur Golu, Turkey. *Threatened Waterfowl Specialist Group News* 4: 7-8.
- Grieve, A. 2001. *White-headed Duck concentration in Kazakhstan*. Threatened Waterfowl Specialist Group Forum. 28 October 2001.
- Heath, M. F. and Evans, M. I. (ed.). 2000. *Important Bird Areas in Europe: Priority sites for conservation*. 2 vols. BirdLife International, Cambridge.

- Kovshar, A. F. 1999. Rare, disappearing and vulnerable birds of Kazakhstan. In *Territorial aspects of birds protection in Middle Asia and Kazakhstan*, pp 77-84. Moscow. (in Russian)
- Meyer de Schauensee, R. 1984. *The Birds of China*. Oxford University Press, Oxford.
- Oliver, D. G. 1919. Spot Bill Duck in Kashmir. *J. Bombay Nat. Hist. Soc.* 26: 675.
- Plowden, W. C. 1887. Letter about Stiff-tailed Duck from Gujrowla, Philibhit. *Stray Feathers* 10: 520–521.
- Singh, G. 1993. *A checklist of birds of Punjab*. Punjab Government Press, Punjab.

Appendix A: List of all sites with White-headed Duck records in the Central Asian region, 1980-2002

Country	Site name and location	Coordinates	Breeding season	Migration period	Non breeding period	Maximum number of birds recorded and year
China	Akto, Near Kashi City, western Xinjiang Autonomous Region	39°10'N, 76°00'E		X		Some specimens during 1981-1988
	Ordos, Inner Mongolia Autonomous Region	39°48'N, 109°20'E		X		1 in 1999
	East Dongting Lake, Hunan Province	29°27'N, 112°48'E			X	1 in 2001
India	Amakhera wetland, Uttar Pradesh	27°31'N, 78°19'E			X	1 in 1997
	Harike Lake, Punjab	31°10'N, 75°00'E		X		1 in 1984
Iran	Alagol Lake, Mazandaran	37°21'N, 54°35'E			X	534 in 2002
	Arjan, Fars Province	29°37'N, 51°59'E			X	37 in 1994
	Bozozjigh, Eastern Azerbaijan	37°27'N 46°46'E	X	X		88 in 1998
	Bur Alan, West Azerbaijan	39°40'N, 44°45'E			X	7 in 2002
	Chaghakhor, Upper Karun River, Zagros Mountains	31°55'N, 50°54'E		X		10 in 1997
	Gandoman, Upper Karun River, Zagros Mountains	31°50'N, 51°07'E		X		6 in 1997
	Ghorigol, Eastern Azerbaijan	37°55'N, 46°42'E	X			108 in 1998
	Gorgan Bay, Mazandaran	36°52'N 53°53'E			X	7 in 1991
	Gorgor, south Iran	Approx 31°30'N, 48°30'E	X			4 in 1997
	Haftbarm, Fars Province	29°40'N, 52°10'E			X	12 in 1993
	Helleh Region	29°09'N, 50°55'E			X	173 in 1988
	Izeh, northeast of Ahwaz, Khuzestan Province	31°54'N, 49°52'E			X	17 in 1996
	Miankaleh Protected Area, southeast Caspian Sea, Mazandaran	36°50'N, 53°45'E			X	50 in 2002
	Parishan Lake, Zagros Mountains, Fars Province	29°31'N, 51°48'E			X	52 in 1993
	Tashk and Kamjan, Fars Province	29°40'N, 53°30'E			X	26 in 1999
	Ulma Gol Lake, Mazandaran	37°25'N, 54°38'E			X	1,450 in 1995
	Yanigh, Eastern Azerbaijan,	37°25'N, 46°59'E	X			17 in 1996
Zoulbin, Eastern Azerbaijan,	37°27'N 46°51'E	X			45 in 1998	
Kazakhstan	Alakol Lake, east Kazakhstan	45°59' N, 81°28'E	X			6 in 1998
	Kaldykol and Biyikkol Lakes	43°N, 70°30'E	X			2 in 2001
	Kamysh-Samara Lakes	48°54'-48°57'N, 49°34'-49°42'E	X	X		490 in 186
	Korgalzhyn-Tengiz Lakes Region (Kokai, Kumkol, Kysylkol, Kumdykol, Nygis, Saumalkol, Sholak and Zhumaj Lakes)	50°10'-50°50'N, 68°40'-71°00'E	X	X		4,021 in 2002

Country	Site name and location	Coordinates	Breeding season	Migration period	Non breeding period	Maximum number of birds recorded and year
Kazakhstan	Kyzylkol Lake, south Kazakhstan	43°44'N, 69°30'E	X	X		2,838 in 2001
	Naurzumskiy Lakes	Approx 51°30N, 64°00E	X			230 pairs with other lakes in southern Russia and northern Kazakhstan 1969-1989
	Presnovskiy and Mibalykskiy Lakes	approximately 54°N, 70°E	X			230 pairs with other lakes in southern Russia and northern Kazakhstan 1969-1989
	Sorbulak Lake, Almaty, southeast Kazakhstan	43°36'N, 76°47'E		X		6 in 2001
Mongolia	Chono Kharaikh Riverl Delta/ Khar Lake	approximately 48°05'N, 93°10'E	X			2 in 1995
	Hodoo Lake, near Zereg	approximately 47°10'N, 92°50'E	X			1 in 1995
	Khar Us Lake (southern section)	47°45' - 48°23'N 91°57' - 92°49'E	X	X		238 in 1998
	Khayrgas Lake	48°58' - 49°20'N 92°48' - 93°48'E	X			8 in 1981
	Olon Lake, Bulgan	49°52'N, 102°38'E	X			5 in 2001
	Uvs Lake (Shuvuun Tsuglaan Lake, west Uvs Lake, 50°12'N, 92°10'E; Jiree and Tes river mouth, east Uvs Lake 50°25'N, 93°05'E)	49°59' - 50°41'N 92°13' - 93°25'E	X			"Large colony" in 2000
Pakistan	Jahlar Lake	32° 29'N, 72°07'E			X	132 in 1988
	Khabekki Lake	32° 37'N, 72° 14'E			X	734 in 1983
	Kharal Lake	30°53'N, 73°35'E			X	48 in 1987
	Ucchali Lake	32° 33'N, 72° 01'E			X	620 in 1987
Russia	Ataich'ie Lake, Omsk Region	54°30'N, 75°40'E	X			4 in 1997
	Atkul Lake, Chelyabinsk Region		X			2 in 1992
	Baganskiy Lakes, southwest Siberia	54°09'N, 78°23'E	X			230 pairs with other lakes in southern Russia and northern Kazakhstan 1969-1989
	Belye Lake, Krasnoyarsk Region	54°40'N, 90°10'E	X			20 in 1985
	Butash Lake, Chelyabinsk Region		X			3 in 1994

Country	Site name and location	Coordinates	Breeding season	Migration period	Non breeding period	Maximum number of birds recorded and year
Russia	Chany Lakes, southwest Siberia	54°52'N, 77°27'E	X			230 pairs with other lakes in southern Russia and northern Kazakhstan 1969-1989
	Chebarkul Lake, Ural, Chelyabinsk Region	55°07'N, 60°35'E	X			Recorded in 1995 and 1997
	Chervyanoe Lake, Sverdlovsk Region		X			Observed in 1995
	Dengino Lake, Chelyabinsk Region		X			2 in 1996
	Duvankul Lake, Chelyabinsk Region	54°40'N, 61°30'E	X			2 in 1994
	Fadiha Lake, Zdvinskij District, Novosibirsk Region	54°36'N, 78°12'E	X			1 in 1997
	Kartabyz Lake, Chelyabinsk Region	54°30'N, 62°30'E	X			1 in 1994
	Katai Lake, Ural, Chelyabinsk Region	55°04'N, 62°05'E	X			1 in 1992
	Kurlady Lake of Ural, Chelyabinsk Region	55°01'N, 61°05'E	X			20 in 1980-2001
	Lake near Cheraskul, Chelyabinsk Region		X			2 in 1987
	M. Bugodak Lake, Chelyabinsk Region		X			1 in 1995
	Malyi Sarykul Lake, Chelyabinsk Region	54°50'N, 61°30'E	X			5 in 1994
	May-Sor Lake, Omsk Region	54°27'N, 75°38'E	X			7 in 2000
	Mayan and Alakul Lakes, Chelyabinsk Region		X			3 in 1994
	Momynkul Lake, Chelyabinsk Region		X			1 in 1987
	Orlovka village Lake, Chistoozernij district, Novosibirsk Region	54°28'N, 76°39'E	X			1 in 2000
	Selezyan Lake, Chelyabinsk Region		X			3 in 1992
	Shira Lake, Krasnoyarsk Region	54°30'N, 90°10'E	X			5 in 1980
	Sykandyk Lake, Chelyabinsk Region		X			7 in 1992
	Sylkino Lake, Omsk Region		X			4 in 1998
	Tishki Lake, Chelyabinsk Region	55°50'N, 61°40'E	X			3 in 1993
	Treustan Lake, Chelyabinsk Region		X			4 in 1992
	Travyanoe Lake, Chelyabinsk Region		X			1 in 1980s

Country	Site name and location	Coordinates	Breeding season	Migration period	Non breeding period	Maximum number of birds recorded and year
Russia	Uelgi Lake, Chelyabinsk Region	55°40'N, 61°30'E	X			2 in 1995
	Ulianovskoe Lake, Baganskij district, Novosibirsk Region	54°08'N, 78°11'E	X			10 in 1998
	Utich'ie Lake, Chelyabinsk Region		X			1 in 1994
	Zabaluevo Lake, Ural, Chelyabinsk Region	54°06'N, 63°00'E	X			6 in 1990-1995
Turkmenistan	Balkhan Bay, Caspian Sea	40°01'N , 52°48'E		X		127 in 1993
	Becovichbay, southeast Caspian Sea	39°32'N , 52°50'E			X	64 in 1984
	Krasnovodsky Bay, southeast Caspian Sea	39°42'-40°02'N, 52°53' -53°32'E	X	X	X	313 in 1994
	Kianly Bay, southeast Caspian Sea	40°12'N , 52°44'E			X	7 in 1999
	Lakes/rivers of mainland Turkmenistan				X	636 in 1986
	Sarakamysh Lake	41°50'-42°20'N , 56°52'-57°50'E		X		930 in 1980
	Southeast Caspian Sea, Between Carabogasgol – Gasankuly (include Balkhan Bay, Becovichbay, Krasnovodsky Bay, Kianly Bay and Severo-Chelecensky Bay)	37°23'-41°08'N , 52°60' -53°59'E			X	820 in 1998
	Severo-Chelecensky Bay, southeast Caspian Sea	39°33'-39°42'N, 53°14'-53°30'E			X	223 in 1991 with around area
	Sutandag Lake, southern Turkmenistan		X			19 pairs in 1989
Uzbekistan	Akushpa Lake, Sudochye Wetlands	58°30'N, 43°14'E	X	X		More than 3,000 in 1999
	Aydar Lake	40°53'N, 66°35'E		X	X	Several tens in 1997-2002
	Dengizkul Lake, south Uzbekistan, near border with Turkmenistan	39°07'N, 64°10'E			X	185 in 2000
	Deukhona lake	39°12'N, 64°39'E			X	6 in 2002
	Karateren Lake, Sudochye Wetlands	58°30'N, 43°14'E		X		9 in 2001
	Rogatoe Lake, near Uchkuduk Town, central Kyzylkum Desert	42°02'N, 63°37'E	X			1 in 1982
	Tuzkan Lake	40°37'N, 67°11'E				5 in 2001
	Tudakul Lake	39°52'N, 64°46'E		X		2 in 2000

Note: X indicate the records of the White-headed Duck in different period.

Appendix B: Site information sheets for key sites/areas for the White-headed Duck in Central Asian region.

- Notes: 1. Questionnaires disseminated to experts in the region as part of the project have solicited detailed site information on key sites/areas. The attached sheets summarise the information received.
2. Information for selected sites for which information was accessible in the literature have been prepared by the authors.

Information Sheet 1. Important sites for the White-headed Duck in China

Site Name	Wetlands of Akto
Area (ha)	Not available
Location	Kizilsu Kirgiz Autonomous Prefecture, western Xinjiang Autonomous Region. Near Kashi City and Artux City. 39°10'N, 76°00'E, altitude 1200m
Site Description	Arid area with high mountains with rivers, lakes and other wetlands.
Description and intensity of human use of this wetland	High intensity of agriculture and fishing; Increasing animal grazing in and around wetlands; Increasing human population; Hunting.
Protection/ownership of site	No protection.
Current threats and problems	Hunting, high intensity of agriculture, fishing and overgrazing in and around the wetlands. Increasing human population.
Suggestions for conservation of site	Survey and research activities to identify important sites for White-headed Duck. Establish Nature Reserves.
Name of Compiler	Ma Ming

Information Sheet 2. Important sites for the White-headed Duck in India

Site Name	Amakhera Wetland
Area (ha)	500
Location	27°31'N, 78°19'E. 40km from Aligarh, Uttar Pradesh
Site Description	Amakhera is a rain deficient, shallow, slightly saline open water wetland, which is surrounded by agricultural fields and a village on one side. There are many such wetlands within a 30km ² radius of this area, which have large concentrations of waterbirds in winter. Amakhera is always full of thousands of waterbirds, including the Comb Duck, Ruddy Shelduck, Lesser Whistling Duck, Bar-headed Goose, Greylag Goose, Gadwall, Pintail, Spot-billed Duck, Wigeon, Garganey, Shoveller, and Common Pochard.
Description and intensity of human use of this wetland	Amakhera wetland is surrounded by agricultural fields and a village on one side. This wetland is used by villagers for bathing, washing clothes and as a source of drinking water for the livestock.
Protection/ownership of site	The wetland is protected by the villagers and hunting is strictly prohibited. It is owned by the Revenue Department.
Current threats and problems	May be drained for agriculture in the future.
Suggestions for conservation of site	The site should be surveyed and monitored regularly, especially during the winter season, when migratory birds are present.
Name of Compiler	M. Zafar-ul Islam

Information Sheet 3. Important sites for the White-headed Duck in Iran

Site Name	Alagol Lake and Ulmagol Lake
Area (ha)	1,180 (Alagol 900 ha and Ulmagol 280 ha)
Location	37°21'N, 54°35'E. The lakes, Alagol and Ulmagol are situated in the Province of Mazandaran, on the Turkoman steppes near the border with Turkmenistan. The lakes are situated about 60km southwest from the town of Gorgan.
Site Description	<p>The site comprises a group of small lakes with associated marshes on the rolling grassy steppes to the east of the Caspian Sea. Alagol Lake is a slightly saline, seasonal lake. It lies about 6km southwest of Ulmagol Lake. The Ulmagol Lake is a freshwater lake. It supports reedbeds, and some aquatic vegetation. It is subject to wide fluctuations in water level, and occasionally dries out completely.</p> <p>The Alagol and Ulmagol lakes are good representative examples of natural saline and freshwater lakes characteristic of the vast plains to the east of the Caspian Sea. The lakes provide wintering habitat for four threatened species of birds: Dalmatian Pelican, Lesser White-fronted Goose, Imperial Eagle and White-headed Duck. Together they regularly support over 20,000 waterbirds in winter, and over 1% of the regional Middle East breeding populations of Gadwall, Tufted Duck and Common Coot.</p> <p>The lakes are utilised by a wide variety of waterbirds during the migration season and in winter. They are especially important for Greater Flamingo, Greylag Goose, Pochard, Smew and Common Coot. Breeding species include Great Crested Grebe, Black-winged Stilt, Kentish Plover, Slender-billed Gull and Eurasian Penduline Tit. Black Stork has been recorded in summer and may breed.</p>
Description and intensity of human use of this wetland	Activities at the three lakes include grazing, reed cutting, recreational fishing and hunting of waterbirds. The water of Alagol Lake is being used for irrigation by local farmers, and for a fish hatchery. Research on waterbirds has been carried out by the Ornithology Unit of the Department of Environment. They have carried out annual mid-winter censuses since 1969, and breeding surveys on several other occasions. In the early 1990s two MSc students from Teheran University conducted research on the avifauna and physico-chemical characteristics of the lakes. In the surrounding area there are a few small settlements where some agriculture is practised.
Protection/ownership of site	The site has no other designation other than being declared a Ramsar Site in June 1975, and is not legally protected. The site has been listed on the Montreux Record since June 1993 due to problems at Alagol Lake.
Current threats and problems	The Ulmagol Lake is subject to high levels of disturbance from waterbird hunters. At Alagol Lake there is also some disturbance from hunting, but on a much smaller scale since Alagol Lake is less accessible. The extraction of water from Alagol Lake for irrigation purposes and for a fish hatchery has resulted in lower water levels at the lake, especially in summer.
Suggestions for conservation of site	None
Name of Compiler	David Li Zuo Wei, based on the Ramsar Information Sheet

Site Name	Ghorigol, Zoulbin, Yanigh and Bozjigh, Eastern Azerbaijan
Area (ha)	Ghorigol 200 ha, Zoulbin 10 ha, Yanigh 4 ha, Bozjigh 10 ha
Location	Ghorigol, 37°55'N 46°42'E, Eastern Azerbaijan, 40km west to Tabriz, near to Bostan Abad Zoulbin, 37°27'N 46°51'E, Eastern Azerbaijan, 24km to Hashtroud Yanigh, 37°25'N 46°59'E, Eastern Azerbaijan, 35km to Hashtroud

Site Name	Ghorigol, Zoulbin, Yanigh and Bozjigh, Eastern Azerbaijan
	Bozjigh, 37°27'N 46°46'E, Eastern Azerbaijan, 40km to Hashtrud
Site Description	Ghorigol, Altitude 1992m, permanent wetland, western and southern areas are covered by reed beds, average depth: 2m, frozen five months of the year. Zoulbin, Altitude: 1650m, permanent wetland, in drought conditions a major part of this wetland will be dry, frozen five months of the year. Yanigh, Altitude: 1650m, permanent wetland, in drought conditions a major part of this wetland will dry up Bozjigh, Altitude: 1650m, permanent wetland, in drought condition a major part of this wetland will dry up, frozen for four months of the year.
Description and intensity of human use of this wetland	Recreational, agricultural and grazing area
Protection/ownership of site	Ghorigol, Governmental, Non Hunting Area, Ramsar Site Zoulbin, No conservation status, public land
Current threats and problems	Overgrazing, over-extraction of water, drought, shooting
Suggestions for conservation of site	Upgrade status to Wildlife Refuge
Name of Compilers	Hamid Amini, Sadegh Sadeghi Zadegan and Yavar Shahbazi

Site Name	Miankaleh Protected Area and Gorgan Bay
Area (ha)	97,200 ha
Location	36°50'N, 53°17'E. Miankaleh Peninsula and Gorgan Bay are situated in the Province of Mazandaran, in northern Iran. They are located at the southeast extremity of the Caspian Sea
Site Description	Gorgan Bay is a shallow, brackish embayment, almost cut off from the Caspian Sea by the 60km long Miankaleh Peninsula, a low, sandy peninsula with coastal dunes, pomegranate scrub and grassland. There are extensive freshwater marshes and seasonally flooded <i>Tamarix</i> woodland at the west end of the bay, and marshes along its south shore. The wetlands of Miankaleh Peninsula and Gorgan Bay form an outstanding example of a natural sand spit/coastal lagoon system characteristic of the south Caspian. They play a substantial hydrological and ecological role in the functioning of the coastal systems of the southeast Caspian. Miankaleh wildlife refuge is one of the finest waterbird reserves in the Western Palearctic region. The wetlands provide wintering habitat for four species of globally threatened birds: Dalmatian Pelican, Pygmy Cormorant, White-headed Duck and Imperial Eagle. Gorgan Bay forms important spawning and nursery grounds for various fish species. Most of Miankaleh Peninsula is covered with a carpet of herbaceous plants, and grasses such as <i>Agropyron</i> , <i>Bromus</i> , <i>Dactylis</i> , <i>Cynodon</i> and <i>Festuca</i> . The western half also supports shrubby woodlands with scattered pomegranate, hawthorn, <i>Rhamnus</i> and blackberry. There are a few large willow trees planted around shepherds' houses. Much of the shoreline of the bay is fringed with a broad belt of rush and there are some large areas of glasswort flats. The marshes at the west end of the bay are dominated by sedges, with small patches of reed-beds, clumps of rush and a large stand of tamarisk. The marshes around Gorgan Bay are eutrophic due to the inflow of numerous streams, agricultural run-off and irrigation channels. The reserve is very important for its large population of raptors. Twenty-eight species have been recorded. Breeding species include Osprey, Short-toed Eagle and White-tailed Eagle. A large number of passerines remain in winter. At least

Site Name	Miankaleh Protected Area and Gorgan Bay
	288 species of birds have been recorded. At Miankaleh Peninsula and Gorgan Bay the Golden Jackal and Wild Boar are abundant in the reserve, and the Jungle Cat also occurs. Caspian Seals occasionally come to the Caspian shore.
Description and intensity of human use of this wetland	Gorgan Bay and the adjacent inshore waters of the Caspian Sea support an important commercial fishery. At Miankaleh Peninsula there is a fish processing factory, some farms, cultivation of cotton and wheat and small villages. Grazing by domestic livestock such as sheep, water buffaloes, goats, cows and horses.
Protection/ownership of site	The entire area of Miankaleh Peninsula and Gorgan Bay was designated as a protected region in May 1970. The wildlife refuge, to which 68,800 ha of the area was designated between 1970 and 1975, was designated as a Ramsar Site in June 1975 and a UNESCO Biosphere Reserve in June 1976.
Current threats and problems	Poaching and over-grazing by domestic livestock are relatively minor threats at Miankaleh Peninsula and Gorgan Bay. Irrigation schemes on agricultural land to the west and the south reduce the flow of freshwater into the marshes and the bay, especially in summer. The major threat to the site is the construction of an asphalt highway down to the centre of the peninsula to provide easy access to the fishery stations along the beach. The greatly increased access to the peninsula will inevitably lead to increased pressure for settlement, increased farming activities and increased poaching.
Suggestions for conservation of site	None
Name of Compiler	David Li Zuo Wei, based on the Ramsar Information Sheet

Site Name	Parishan Lake and Dasht-i Arjan, Fars Province
Area (ha)	Total area: 6,200 ha; Parishan Lake, 4,000 ha; Dasht-i Arjan, 2,200ha
Location	29°30'N, 52°00'E. Parishan Lake and Dasht-e-Arjan are situated in the Province of Fars, in the Zagros mountains, 40-80km west of Shiraz and 15-25km south of Kazerun.
Site Description	Elevation: 2,000 m above sea level (ASL) (Dasht-i Arjan); 853 m ASL (Parishan Lake). The wetland of Dasht-e-Arjan is an outstanding example of a freshwater wetland characteristic of the highlands of western Iran. Parishan Lake is a good example of a saline wetland characteristic of the same highlands. They support five species of threatened birds: Dalmatian Pelican, Marbled Teal, Ferruginous Duck, White-headed Duck and Imperial Eagle. Both wetlands support a very diverse flora and fauna, and thus maintain the genetic and ecological diversity of the region. In winter, the lakes hold over 20,000 waterbirds. They also support over 1% of the regional wintering populations of White Pelican, Greater Flamingo, 11 species of ducks, Common Coot, Eurasian Crane and Black-headed Gull. During breeding season, large breeding colonies of herons and ibises can be found at the lakes, as well as over 1% of the regional populations for Glossy Ibis and Eurasian Spoonbill.
Description and intensity of human use of this wetland	There are subsistence fishing, reed-cutting and extensive grazing by domestic livestock. The Ornithology Unit of the Department of the Environment has carried out annual mid-winter censuses since 1967. There are plans to build a visitor centre. This site is renowned for its spectacular scenery. In the surrounding area there are a few small settlements with orchards and gardens, some wheat cultivation and other crops.
Protection/ownership of site	A national park of 65,750 ha was established in 1972, but at the end of the 1970s after the revolution it was downgraded to the Arjan Protected Area of 52,800 ha. The site was designated as a Ramsar site on 23 June 1975. The area of the original national park was designated an UNESCO Biosphere Reserve in June

Site Name	Parishan Lake and Dasht-i Arjan, Fars Province
	1976. There is a game guard station on a peninsula overlooking the western part of Parishan Lake. At both lakes hunting has been prohibited since 1973. The Ramsar Management Guidance Procedure Mission that visited the lakes in 1992 made several recommendations. The most important are to demarcate the borders of the site clearly with signs, and to prohibit any further drainage activities at Parishan Lake.
Current threats and problems	Parishan Lake is under considerable threat from various sources, while Dasht-e-Arjan remains in reasonably good condition. Some 20 hectares of marsh at the extreme northwest corner of Parishan Lake were drained for agriculture about 20 years ago. Elsewhere around this lake, wet meadows have been replaced by cultivated fields. A small area of fish ponds was established on the plains to the west of the lake in the early 1980s, and it is reported that three species of carp have been introduced to the lake. There has been a considerable increase in fishing activities, and the widespread use of outboard motor boats instead of traditional reedboats has resulted in disturbance to the waterbird populations. Poaching remains a problem, as well as the accidental killing of waterbirds in fishing nets. At Dasht-e-Arjan poaching is also a problem. Two sets of high-tension power lines that cross the lake are dangerous to birds and are spoiling the beautiful scenery of the lake. One of the lines has not been in use since it was constructed in the late 1970s.
Suggestions for conservation of site	None
Name of Compiler	David Li Zuo Wei, based on the Ramsar Sheet information Sheet

Information Sheet 4. Important sites for the White-headed Duck in Kazakhstan

Site Name	Kyzylkol Lake, Dzhambul region, south Kazakhstan
Area (ha)	500-750 ha
Location	165km north-west of Dzhambul, south Kazakhstan. 43°44'N, 69°30'E.
Site Description	A saline, shallow lake with submerged vegetation but no emergent or marginal vegetation. It is surrounded by salt-marsh plants set back from the water edge. There is no outflow and only one freshwater inflow at the southeastern corner. The lake is situated in a semi arid desert/steppe area. Up to 200,000 waterbirds (coot, ducks and waders) are known to occur in autumn.
Description and intensity of human use of this wetland	Little human use, perhaps some light grazing. About a dozen shooting butts are found on shore around the lake probably used by 3 – 4 hunters several times a week, but there is no evidence of heavy shooting using boats. When shooting was observed, the birds moved out of range into the middle of the lake.
Protection/ownership of site	There is no official, legal protection for this site. Ownership details are not known.
Current threats and problems	Current threats are not known. There is intensive recreational use (shooting) in summer but not during periods when it is used by waterbirds. Over-hunting could become a problem in the future.
Suggestions for conservation of site	Site should be declared a Protected Area and hunting reduced/restricted or stopped.
Name of Compiler	Andrew Grieve

Site Name	Tengiz-Korgalzhyn Lakes Region
Area (ha)	1,200,000 ha
Location	The lakes are situated in the Akmola District (Akmolinskaya Oblast), northeast of the Aral Sea in central Kazakhstan, 50°10'-50°50'N, 68°40'-71°00'E. Korgalzhynskij Zapovednik, Akmolinskaya Region, 120km southwest of the capital Astana.
Site Description	<p>Tengiz-Korgalzhyn Region consists of a complex system of fresh to saline water with many lakes separated by extensive reed beds and with a network of deep water channels with varying degrees of salinity. The whole system is fed by the Nura River which ends in the Tengiz Lake. Korgalzhyn Lake is separated by a series of dams from the huge and saline water body of the Tengiz Lake. Tengiz Lake is a saline lake of 156,000 ha. The water levels of the lakes depend on the inflow from the Nura and Kulanutpes rivers, the operation of sluices on the dams across the rivers, the amount of flooding, and the evaporation rates. The lake water has a variety mineral composition, particularly high in sulphates and chlorides. Korgalzhyn Lake is a lake of 39,600 ha with varying degrees of salinity. It comprises a network of deep water channels (0.5-2.5 m) supports rich aquatic vegetation. The average temperature in January is -17°C and in June it is 20°C. The number of days with an average temperature below zero is 150 days per year.</p> <p>The vegetation at and around Korgalzhyn Lake comprises reed beds with <i>Phragmites communis</i>. At the lakeside the grasses <i>Stipa lessingiana</i>, <i>Agropyron repens</i>, <i>Bromus inermis</i>, <i>Festuca sulcata</i> and <i>Calamagrostis epigeios</i> grow, as well as the flowering plants <i>Pyrethrum achilleifolium</i>, <i>Spiraea hypericifolia</i>, <i>Artemisia</i> spp., <i>Halocnemum strobilaceum</i> and <i>Atriplex cana</i>. Korgalzhyn Lake also supports large zooplankton and phytoplankton and fish populations. Korgalzhyn and Tengiz Lakes are good examples of saline lakes characteristic of the northern Kazakhstan.</p> <p>In the vicinity of the Nature Reserve (mainly northeast, east and southeast) is a large number of other lakes situated mainly around the little town of Korgalzhyn, possessing varied hydrologic regimes with saline or fresh water. The most important ones for White-headed Ducks are: Saumalkol Lake, Zhumaj Lake, Kumkol Lake, Kumdykol Lake and Nygis Lake and others (all with saline water). These lakes usually have considerably less vegetation than the lakes inside the reserve. Their importance is probably less as breeding sites, but more as staging areas.</p> <p>Peak numbers of White-headed Ducks for the area, i.e. Korgalzhyn Lake and adjacent lakes represent about 4,000 birds during September while the local breeding population is estimated at probably a few hundred pairs. The lakes are of great importance as breeding, moulting and especially resting stations for an enormous number of Anatidae and waders of many different species during their migration. The wetland is the main nesting site for 5,000-18,000 pairs of Greater Flamingo (Tengiz Lake). Dalmatian Pelicans breed with up to 500 pairs at Korgalzhyn Lake. Lesser White-fronted Geese occur with some hundreds and Red-breasted Geese with some thousands of individuals in the area and adjacent territory during migration. Other non-breeding (summer) visitors in high numbers include Common Shelduck and Ruddy Shelduck. The site is a moulting refuge for numerous non-breeding birds including European Wigeon, Northern Pintail, Gadwall, Common Pochard and Greylag Goose.</p>
Description and intensity of human use of this wetland	In the area around Korgalzhyn Lake all activities are prohibited except research. In the area around Tengiz Lake there occurs some hunting, hay cultivation and some exploitation of natural resources. Studies have been conducted on the ecosystem and on waterbird populations.

Site Name	Tengiz-Korgalzhyn Lakes Region
Protection/ownership of site	The site was declared a Zapovednik (strictly protected nature reserve) in 1968. It was also a Ramsar site during Soviet Union times but Kazakhstan has not ratified the Convention since it became independent.
Current threats and problems	Favourable conditions for the White-headed Duck depend a lot on the water regime and on water management. Strong natural fluctuations in the water regime are typical for the Central Asian region. The water management is weak and function and control of dams are not effective. The impact of contaminants (mercury for instance) needs to be evaluated, particularly in the Nura River.
Suggestions for conservation of site	The site is legally protected but support in conservation management is necessary.
Name of Compiler	Goetz Eichhorn/Lars Lachmann

Information Sheet 5. Important Sites for the White-headed Duck in Mongolia

Site Name	Khar Lake
Area (ha)	50,000 ha
Location	Located to the east of Har Us Lake in the valley of Great Lakes, Khovd Aimag (Province). 47°58' - 48°13'N, 93°00' - 93°25'E. Altitude 1106m ASL.
Site Description	A large freshwater lake with a maximum depth of 7m. It is connected to Khar Us Lake in the west, and its outflow flows into Dorgon Lake, immediately to the southeast. The lake is also connected to the Dzavkhan River. The faunal and floral characteristics are very similar to those of Khar Us Lake. The area has an extreme continental climate modified by altitude. The annual precipitation is about 300mm. A substantial area of the lake is occupied by macrophytes, mainly <i>Potamogeton</i> and <i>Nymphaea</i> sp.
Description and intensity of human use of this wetland	Livestock grazing.
Protection/ownership of site	The area is included in Khar Us lake National Park. The site is state owned. This site was nominated as a Ramsar site jointly with Khar Lake and Dorgon Lake in April 1999
Name of compiler	David Li Zuo Wei, based on Scott 1989

Site Name	Khar Us Lake
Area (ha)	150,000 ha
Location	47°45' - 48°23'N, 91°57' - 92°49'E, in the Valley of Great Lakes close to the Altai Mountains. 180km south of the Ulaangom City and 25km east of the Khovd City, Khovd Aimag, western Mongolia.
Site Description	This is a fresh water tectonic lake located at 1,157 m ASL, near ancient ruins 20km to the east of the Aimag/province centre. This is one of four lakes with an area of over 1,000km ² . Its area, including islands, is 1,852km ² with 72km length, 17km width and a 306.8km shoreline perimeter. It is fed by water from a catchment of about 70,450km ² . Khovd, Buyan and Tsenkher Rivers flow into the lake and only Chono Kharaiikh River flows out from the lake. Of the 120 islands in the lake, the largest one is Agbashi island with an area of over 400km ² , a length of 30km and it rises 272 metres above the lake. This island divides the lake into two parts: the Northern and Southern "oceans". The two "oceans" are connected through the straits of Lun and Yum. The Northern "ocean" is 400km ² in area with a depth of 2 metres. It is covered with reeds. The depth of the Southern "ocean" reaches 4.5 meters. Water temperatures during the period of

Site Name	Khar Us Lake
	<p>June to August is between 20⁰C and 23.5⁰C, whereas in the winter season it falls to 1⁰C -1.5⁰C at the surface and 5.1⁰C -6.5⁰C on the bottom. In the period between November and April it is covered with 0.8-1.2 meters of ice. Water transparency is 0.7-1.0 metres. The bed is covered with thick clays and smells of hydrogen sulphide. Along some of the shores of the lake there are sands and thick growths of water plants. Up to 80% of the lake is covered with macrophytes; the dominant forms are floating-leaved such as <i>Potamogeton</i> sp. and <i>Nymphaea</i> sp. Total mineral content is 348.03-242.08 mg/l. The pH varies between 7.7 and 7.83.</p> <p>Vast reed beds and extensive aquatic plant communities provide a suitable habitat for a large number of breeding and migratory waterbirds, including the globally threatened Swan Goose, Ferruginous Duck, White-headed Duck and Relict Gull. Three species of fish endemic to Western Mongolia (<i>Oreoleuciscus</i> sp.) occur in these lakes.</p>
Description and intensity of human use of this wetland	Human use includes fishing, harvesting of muskrats and livestock grazing. The Muskrat was introduced into the lake area in 1967, and has become very abundant. Some 5,000 individuals were harvested annually in the 1970s. The lake supports a locally important fishery. Since it was protected as a National Park in 1997, muskrat harvesting and fishing are now prohibited.
Protection/ ownership of site	The area is included in the Khar Us Lake National Park. It is state owned. This site was nominated as a joint Ramsar site together with Har Lake and Dorgon Lake in April 1999.
Current threats and problems	The habitat for birds is getting worse due to livestock grazing, reed cutting, steppe fires in spring and autumn, and illegal hunting. The proposed construction of a hydroelectric power plant adjacent to the national park, overgrazing of pastureland and the introduction of the Muskrat are the main threats to the ecological integrity of the site.
Suggestions for conservation of site	None
Name of compilers	N. Tsevenmyadag and S. Gombobaatar

Site Name	Khyargas Lake
Area (ha)	140,700 ha
Location	Located in the valley of Great Lakes in western Mongolia, 100km southwest of Ulaangom, Khovd Aimag. 48°58'- 49°20'N, 92°48'-93°48'E, Altitude 1029m ASL.
Site Description	It is one of the biggest saline lakes in Mongolia, in the lowest depression in the Valley of Great Lakes between the Altai and Khanggai mountain ranges. The lake is 75km long and up to 31km wide, and has a maximum depth of 80m. It is connected to a nearby freshwater lake, Airag Lake. The lake has ice cover from October to May. The climatic condition is extreme continental modified by altitude. The annual precipitation is about 300mm.
Description and intensity of human use of this wetland	Some livestock grazing
Protection/ ownership of site	The area is included in Khyargas Lake National Park. State owned.
Current threats and problems	Not available
Suggestions for conservation of site	None
Name of compiler	David Li Zuo Wei, based on Scott 1989

Site Name	Olon Lake
Area (ha)	500 ha
Location	49°52'N, 102°38'E, in the Basin of Egiin Gol River, 2-5km south of Teshig soum Village and 140km northwest of Bulgan city, Bulgan Aimag, central Mongolia.
Site Description	This is a freshwater tectonic lake located at 980 metres ASL. This is one of 20 lakes in the area. It has an area of 0.5km ² . with a length of 0.9km and a width of 0.7km. The shoreline perimeter is 2.2km. It is connected to the Egi River. Water temperature is 18.9°C-20.1°C in June and August. The bottom of the lake is covered with slime. There are a lot of water plants growing in the lake. The lake has an abundance of fish and migratory birds. Muskrats have been introduced into the lake area.
Description and intensity of human use of this wetland	Fishing, harvesting of Muskrats and livestock grazing.
Protection/ownership of site	No protection, state owned.
Current threats and problems	The reeds are decreasing due to the introduction of the Muskrat.
Suggestions for conservation of site	None
Name of compiler	N. Tsevenmyadag

Site Name	Shuvuun Tsuglaan Lake on the west side of Uvs Lake
Area (ha)	400 ha
Location	In Sagil soum, Uvs Aimag, western Mongolia. About 23km northeast of the city of Ulaangom and 10km east of Uvs Lake. 50°10'-50°13'N, 92°08'-93°12'E.
Site Description	This small lake is included in the Uvs Lake basin. It has a length of 2km, a width of 2km and an average depth of 2km. The lake is separated from the Valley of Great Lakes by the Khan-Khukhii ridge. It receives water from the west from the small rivers Nariin and Khondlon. In summer, the water temperature exhibits a gradient from 25°C at the surface to 19°C at the bottom. There is ice cover from October to May. The area has an extreme continental climate.
Description and intensity of human use of this wetland	Local people do not use the area for fishing, agriculture or water supply. The lake and its surrounding area are used for grazing cows, horses, sheep, and goats in summer. No disturbance of waterbirds has been recorded.
Protection/ownership of site	The area is included in the buffer zone of Uvs Lake Strictly Protected Area. State owned.
Current threats and problems	Steppe fire and reed cutting
Suggestions for conservation of site	None
Name of compilers	N. Tsevenmyadag and S. Gombobaatar

Site Name	Tes and Jiree rivers Delta, Northeast site of Uvs Lake
Area (ha)	15,000 ha
Location	Tes soum of the Uvs Aimag in western Mongolia. The area is about 90km northeast from city Ulaangom and about 30km from the village of Tes soum. It is close to the Russian border. 50°20' - 50°31'N, 92°58'-93°17'E. Altitude 759m ASL.

Site Name	Tes and Jiree rivers Delta, Northeast site of Uvs Lake
Site Description	Uvs Lake is the largest in Mongolia in terms of surface area, with a length of 84km, a width of 79km and an average depth of 6km. The lake is separated from the Valley of Great Lakes by the Khan-Khukhii ridge. It has no outlet and has a very large catchment area, receiving water from the east in the Baruntura, Nariyn and Tes rivers. The latter river forms a vast area of marsh to the northeast of the lake, straddling the Russian border. The water is markedly saline (18.8 ppt.), with sulphate and sodium ions being the most important constituents. In summer, the water temperature exhibits a gradient from 25°C at the surface to 19°C at the bottom. There is ice cover from October to May. The area has an extreme continental climate.
Description and intensity of human use of this wetland	Local people do not use the area for fishing, agriculture and water supply. The lake and its surroundings are used for cow, horse, sheep, and goat grazing in the summer, but causes little disturbance for waterbirds.
Protection/ownership of site	The area is included in Uvs Lake Strictly Protected Area. It is state owned. There are 4 different categories of Protected areas in Mongolia. The strongest protection measures are applied in Strictly Protected Areas.
Current threats and problems	Steppe fire and reed cutting
Suggestions for conservation of site	None
Name of compilers	N. Tsevenmyadag and S. Gombobaatar

Site Name	Zavkhan River Delta and Airag Lake
Area (ha)	14,330 ha
Location	In Dzavkhan soum of the Uvs Aimag in western Mongolia. 150km southeast of Ulaangom, and about 20km northeast of the village of Dzavkhan soum. 48°50'-48°57'N, 93°16'-93°32'E.
Site Description	<p>This is a freshwater lake which is located at 1030 m ASL not far from Khyargas Lake. It has an area of 143.3km² with 16km long and 13km wide. The shoreline perimeter is 54km. Its water depth is about 10 m in Galbiin Gobi and 4-8 m in other parts. The water volume is about 820 million cu metres. The shore is steppe-like. In the north, the Galbiin hills separate it from the Khyargas Lake. Where the inflows meet the lake, the soil becomes marshy with densely growing reeds. Zavkhan and Khungui Rivers flow into the lake and it is connected to the wide valley of Galba. Water temperature in July and August is 19.5°C -21.3°C and 2°C -2.5°C at the bottom in February and March. It has ice coverage in the period between November to April and its thickness reaches 1-1.3 metres. The lakebed is composed of sand and whetstone. In addition, there is also black and brown clay which smells slightly of hydrogen sulphide. The mineral content is low (502.02 mg/l) with high concentrations of carbonate and sodium ions. The pH is 8.58.</p> <p>It is an exceptionally important breeding and resting site for a variety of waterbirds and it is the only remaining place in Mongolia where the Dalmatian Pelican regularly breeds. Other noteworthy waterbird species include the globally threatened Swan Goose and the Relict Gull.</p>
Description and intensity of human use of this wetland	Some livestock grazing.
Protection/ownership of site	The area is included in Khyargas lake National Park. The area is state owned. In April 1999, this site was nominated as a Ramsar site.
Current threats and problems	Overgrazing of the pastureland threatens the plant communities around the wetland and disturbs breeding birds.

Site Name	Zavkhan River Delta and Airag Lake
Suggestions for conservation of site	None
Name of compiler	N. Tseveenmyadag

Information Sheet 6. Important Sites for the White-headed Duck in Pakistan

Site Name	Jahlar Lake
Area (ha)	17 ha
Location	32°29'N, 72°07'E; Approx 10km southeast of Uchchali Lake and 10km southwest of Nowshera, Khushab District, Punjab Province.
Site Description	<p>A small saline lake with little marsh vegetation in the Salt Range; similar in general character to the nearby larger Uchchali and Khabekki Lakes. The lake is fed by run-off from the surrounding hills of the Salt Range. The depth varies from 0.2m to 5m depending on the amount of rainfall received. pH values ranged from 9 to 10 in the years of 1989 to 1992.</p> <p>Climatic conditions: Dry subtropical climate with hot summers and cool winters. The annual rainfall varies from 300 mm to 800 mm, and the relative humidity from 22% to 85%.</p> <p>Principal vegetation: The aquatic vegetation includes <i>Carex fedia</i>, <i>Hydrilla verticillata</i>, <i>Juncus sp.</i>, <i>Phragmites australis</i>, <i>Potamogeton crispus</i>, <i>P. pectinatus</i>, <i>Saccharum spontaneum</i>, <i>Typha angustata</i> and <i>Zannichellia palustris</i>. The natural vegetation of the region is a mixture of subtropical semi-evergreen forest and tropical thorn forest with species such as <i>Acacia modesta</i>, <i>Adhatoda vasica</i>, <i>Asparagus gracilis</i>, <i>Cocculus laeba</i>, <i>Cynodon dactylon</i>, <i>Ehretia laevis</i>, <i>Gymnosporia royleana</i>, <i>Olea ferruginea</i>, <i>Reptonia buxifolia</i>, <i>Sageretia lorandehuana</i>, <i>Tamarix aphyllia</i>, <i>Withania coagulans</i>, <i>Zizyphus mauritiana</i> and <i>Z. nummularia</i>.</p> <p>Fauna: Information on the lake is available only after 1987. The lake is probably a regular wintering area for the White-headed Duck (see Table 10 and 11). Other waterbird present in the lake included small numbers of Little Grebe, Black-necked Grebe, up to 601 Common Pochard in January 1990 and up to 112 Common Coot in January 1988. Mammals known to occur in the area include <i>Canis aureus</i>, <i>Vulpes vulpes</i>, <i>Felis libyca</i> and <i>Lepus nigricollis</i>.</p>
Description and intensity of human use of this wetland	<p>The most important use of the lake is aquifer recharge. Agricultural lands around the lake belonging to Jahlar villagers, grazing lands around the lake, and the drinking water supply for the Jahlar village all depend on this ground water. Jahlar is a comparatively small village with a population of about 1,500 and a livestock population of 2,000 head of livestock.</p> <p>The livestock mainly graze in the scrub forests on the hills surrounding the lake. The forests also provide the firewood used by the Jahlar villagers. Some firewood is also taken by the villagers living further away from the lake. Jahlar, before it was declared a wildlife sanctuary, was considered to be a good hunting area and there reportedly used to be hunting of White-headed Ducks but after it was protected, hunting is almost non-existent.</p> <p>There has been no fisheries activity in the lake.</p>
Protection/ownership of site	The lake was declared a wildlife sanctuary in 1993, and all sorts of hunting and habitat disturbance is prohibited. The wetland was also declared (a part of Uchchali Complex wetlands) a Ramsar site on 22 March 1996. The wetland and the surrounding agricultural land are privately owned whereas the hill forests and rangelands are state owned (Government of the Punjab).
Current threats and problems	Hunting activities cause disturbance to waterbird populations. A road has been constructed around two sides of the lake which has effectively stopped the run-off water from entering into the lake. Even though drainage pipes have been put

Site Name	Jahlar Lake
	under the road, these do not effectively drain all the run-off water that remains dammed up behind the road. The water level and the extent of lake, however, depends mainly on the rainfall received during the monsoons which greatly varies from year to year.
Suggestions for conservation of site	<p>Undertaking a study of the status of the wintering population of White-headed Duck on the Salt Range Lakes in northern Pakistan. This should be undertaken in relation to its status on the breeding grounds in Central Asian States to determine the causes of its population decline, and to identify remedial measures for its conservation.</p> <p>Instituting long term monitoring studies of climatic factors and their effect on the physical, chemical and biological characteristics of the Salt Range Lakes.</p> <p>Undertaking a study on the status of the White-headed Duck on the Salt Range Lakes in relation to changes in physical, chemical and biological characteristics of the lakes to suggest appropriate habitat development to ensure the survival of the species.</p> <p>Undertaking a satellite-tracking project to identify the migratory route and stopover sites along the flyway of the White-headed Duck.</p> <p>Strengthening staff capacity of protection agencies at the Salt Range Lakes to prevent illegal hunting of waterbirds which could lead to further population decline.</p> <p>Improving habitat conditions at the Salt Range Lakes, for example, afforestation in surrounding watersheds and restoration of water resources prevent silting of the lakes and further habitat loss.</p> <p>Providing training to local communities in the efficient use of local fuels to minimise the use of vegetation cover at the Salt Range Lakes.</p> <p>Undertaking public awareness raising programmes adjacent villages through organising local Community Based Organisations/Village Organisations (CBOs/VOs) to increase conservation awareness at the Salt Range Lakes.</p> <p>Encouraging development of sustainable ecotourism at the Salt Range Lakes to help improve the local economy.</p> <p>Ensuring management of the Salt Range Lakes, on a Participatory/Benefit sharing basis, with the active involvement of local communities.</p>
Name of compiler	Abdul Aleem Chaudhry

Site Name	Khabekki Lake
Area (ha)	283 ha
Location	32°37' N 72°14'E; 10km northeast of Nowshera and 38km northwest of Khushab, Khushab District, Punjab Province.
Site Description	<p>A shallow, saline lake in the Salt Range, with a little aquatic vegetation but no extensive reed-beds. The lake is fed by local rainfall and several intermittent streams rising in the surrounding hills. The water level in the years 1988-1989 had risen by 30-60 cm, causing an increase in size of the lake and decrease in salinity. Because of subsequent failure of rains, water level again was reduced. The trend remained up to 1992 when due to exceptionally good monsoon rains, the water level rose again. The maximum depth is about 10.5m; a salinity of 5.2 ppt was recorded in April 1987, and a pH of 7.2 in January 1987, 9.5-10 in 1989 and 9 in the years 1991 and 1992. Altitude: 978m ASL. Due to failure of rains in the previous years the extent of the lake has greatly reduced.</p> <p>January 2001: Water level was low due to failure of heavy rains for a long period. Only about one fourth of the lake was covered with water. Waterbird population decreased as a result.</p> <p>January 2002: Due to failure of heavy rains for the last four to five years the extent of the lake has been reduced. Only about 12% of the area is under water.</p>

Site Name	Khabekki Lake
	<p>Very few waterbirds were seen. There were no fisheries activities.</p> <p>April 2002: The extent of lake was further reduced. Only about less than 4 ha under water. There were no fisheries activities.</p> <p>Climatic Conditions: Dry subtropical climate with hot summers and cool winters. The annual rainfall varies from 300 mm to 800 mm, and the relative humidity from 22% to 85%. Temperatures range from an average minimum of 0.5°C in January to an average maximum of 36° C in June.</p> <p>Principal vegetation: The aquatic vegetation includes <i>Carex fedia</i>, <i>Chara sp.</i>, <i>Hydrilla verticillata</i>, <i>Juncus sp.</i>, <i>Najas marina</i>, <i>Phragmites australis</i>, <i>Potamogeton crispus</i>, <i>P. pectinatus</i>, <i>Saccharum spontaneum</i>, <i>Scirpus sp. (short)</i>, <i>Scirpus sp. (tall)</i>, <i>Typha angustata</i>, <i>Vallisneria spiralis</i> and <i>Zannichellia palustris</i>. The natural vegetation of the region is a mixture of subtropical semi-evergreen forest and tropical thorn forest with species such as <i>Acacia modesta</i>, <i>Asparagus gracilis</i>, <i>Cocculus laeba</i>, <i>Cynodon dactylon</i>, <i>Adhatoda vasica</i>, <i>Dodonaea viscosa</i>, <i>Ehretia laenis</i>, <i>Gymnosporia royleana</i>, <i>Olea ferruginea</i>, <i>Rhazya stricta</i>, <i>Sageretia lorandetuana</i>, <i>Reptonia buxifolia</i>, <i>Tamarix aphylla</i>, <i>Withania coagulans</i>, <i>Zizyphus mauritians</i> and <i>Z. nummularia</i>. Most of the natural vegetation around the lake in the valley bottom has been cleared for agriculture, whereas the forest vegetation on the surrounding hills is severely depleted.</p> <p>Fauna: Formerly, and even now, a very important wintering area for the White-headed Duck (see table 10 and 11); and a regular wintering area for Greater Flamingo and many other waterbird. As many as 8,700 waterbirds were recorded in the early 1970s. About 50 bird species are normally recorded during summers. Mammals and amphibians known to occur in the area include <i>Canis aureus</i>, <i>Vulpes vulpes</i>, <i>Felis libyca</i>, <i>Lepus nigricollis</i> and <i>Rana tigrina</i>.</p> <p>Introduced fishes include <i>Labeo rohita</i>, <i>Catla catla</i>, <i>Cirrhinus mrigala</i>, <i>Cyprinus carpio</i> and <i>Ctenopharyngodon idella</i>.</p>
Suggestions for conservation of site	As for Jahlar Lake
Name of compiler	Abdul Aleem Chaudhry

Site Name	Ucchali Lake
Area (ha)	943 ha
Location	32°33'N, 72°01'E; 13km west of Nowshera and 42km northwest of Khushab (District Head quarter) in the Punjab province of Pakistan.
Site Description	<p>A saline lake, the largest in the Salt Range, with little marsh vegetation and almost entirely surrounded by agricultural land. The lake is fed by a small spring, seepage from adjacent irrigated land, and run-off from the surrounding hills of the Salt Range. The water level and salinity fluctuate according to local rainfall. The depth varies from 0.2m to 6m; and water is usually hyper-saline (41.5 ppt in April 1999), and had a pH of 10 in April 1999.</p> <p>Climatic conditions: Dry subtropical climate with hot summers and cool winters. The annual rainfall varies from 300 mm to 800 mm, and the relative humidity from 22% to 85%. Temperatures range from an average minimum of 0.5 °C in January to an average maximum of 36 °C in June.</p> <p>Principal vegetation: Marsh vegetation is confined to a few small patches along the lake shore, but there is a very rich growth of plankton in the lake. The dominant aquatic plants are <i>Carex fedia</i>, <i>Hydrilla verticillata</i>, <i>Juncus sp.</i>, <i>Phragmites australis</i>, <i>Potamogeton crispus</i>, <i>Potamogeton pectinatus</i>, <i>Spergularis marina</i>, <i>Suaeda fruticosa</i>, <i>Haloxylan multiflorum</i>, <i>Cynodon dactylon</i>, <i>Phyla nodiflora</i>, <i>Scirpus sp. (dwarf)</i>, <i>Scirpus sp. (tall)</i>, <i>Paspalum distichum</i>, <i>Typha angustata</i>, <i>Imperata cylindrica</i>, <i>Alhagi camelorum</i>, <i>Rumex sp.</i>,</p>

Site Name	Ucchali Lake
	<p><i>Ranunculus scleratus</i>, <i>Saccharum spontaneum</i>, <i>Typha angustata</i>, <i>Vallisneria spiralis</i> and <i>Zannichellia palustris</i>. The natural vegetation of the region is a mixture of subtropical semi-evergreen forest and tropical thorn forest with species such as <i>Acacia modesta</i>, <i>Adhatoda vasica</i>, <i>Asparagus gracilis</i>, <i>Cocculus laeba</i>, <i>Cynodon dactylon</i>, <i>Dodonaea viscosa</i>, <i>Ehretia laenis</i>, <i>Gymnoporia royleana</i>, <i>Olea ferruginea</i>, <i>Reptonia buxifolia</i>, <i>Rhazya stricta</i>, <i>Sageretia lorandettuana</i>, <i>Tamarix aphylla</i>, <i>Withania coagulans</i> and <i>Zizyphus</i> sp. The natural vegetation around the lake has been cleared for agricultural land.</p> <p>Fauna: A very important wintering area for waterbird, particularly Greater Flamingo 1,360 recorded in January 1987, Anatidae and Common Coot. The number of birds present fluctuates widely from year to year and seems to be critically dependent on the water level and salinity. Between 1,100 and 3,100 Anatidae and Common Coot were reported to winter on the lake in the early 1970s, but much larger numbers have been reported in 1980s, over 100,000 waterbird, largely Common Coot, were present in the winter of 1985/1986, and 50,000 Common Coot were reported in November 1986. It's the most important wintering area for White-headed Duck in Pakistan (see Table 10 and 11). Due to failure of rains and prevention of runoff water from going into the lake in recent years, the area of the lake has greatly been reduced. As a result, the number of waterbirds visiting the lake has greatly declined.</p> <p>January 2001: Due to failure of rains, the water level was very low, and only a few waterbirds were present. As no fresh water is entering into the lake, a pungent smell comes out of the lake and it is difficult to stand near the water. The Flamingo population has shifted from Ucchali to Khabekki lake.</p> <p>January 2002: Due to failure of rains, the extent of the lake has been reduced. As the land under the lake was privately owned, the owners have started cultivating the exposed lakebed. Only a few waterbird were observed in the lake. The deteriorating situation continues. The extent of the lake has been reduced to an area of 16 to 20 ha only (April 2002).</p>
<p>Description and intensity of human use of this wetland</p>	<p>The most important use of Ucchali Lake is for aquifer recharge. Agriculture is the main land use around the lake. Most of the Ucchali village farmland is however, submerged under the lake. Very few farmers have land holdings of more than 4 ha. The average land holding is around 1 ha. The small size of the land holdings and the high productive potential encourages intensive cultivation with several crops per year. More than 80% of the farmed land is located around the lake. The main crops are cauliflower, wheat, maize, jawar, bajra, brassicas, trifolium and coriander. Cauliflower is especially important as the off-season crop that fetches very high prices in the local market.</p> <p>Agriculture is based on proper irrigation and adequate ground water which is amply provided due to the continuously recharged aquifer. Dependence is mainly on well irrigation. Bore-holes have been dug all around the lake, and are a constant drain on the groundwater, resulting in the lake area being further reduced. Additionally, the lake area is being reduced due to failure of rains in recent years.</p> <p>The watershed areas around the lake (small hills from 800 to 1500m ASL) are covered by scrub forests. People collect firewood from the community forests, and also from the Government owned reserved forests, which are relatively dense. As a rough estimate, the village requirement of firewood is around 300 tonnes. People also buy wood from illegal cutters who bring wood from the forests on camel back. An average family with 6-7 members may need around 3 camel loads of firewood per month (cooking gas is also used but only by some families). Community forests are depleting fast, whereas the Government owned forests are also being affected due to illegal cutting of wood. Areas around the lake are also used for grazing livestock. The livestock also graze in the scrub forests in the watershed and cause degradation in the forest cover.</p>

Site Name	Ucchali Lake
	<p>Ugali is the second largest village around the lake (located in the north) with about 3000 inhabitants. Agriculture, however, is not the mainstay of the village population, and only 7% of villagers depend on agriculture. 300 ha area is suitable for agriculture, out of which only 13 ha are irrigated. At present only 16% of land is being cultivated, the rest remains uncultivated, with wild growth which is utilised by livestock. Livestock number around 5000 including goats, sheep, cows, buffaloes, donkeys and camels, but the interest in rearing livestock is decreasing due to paucity of grazing lands around the village and distantly located scrub forests.</p> <p>The wetland has been declared a Game Reserve but special shooting permits have never been issued. Yet there is a history of illegal shooting on the lake. Limited nature-oriented recreation activities are carried out. Due to hypersalinity, fisheries introduction has not been successful, hence this activity is non-existent. Due to marshy conditions on the periphery of the lake some tall grasses grow which are harvested for thatching purposes.</p>
Protection/ ownership of site	The wetland was declared a Game Reserve in 1985. Limited hunting is allowed in a Game Reserve on special permit basis, only two permits in one shooting season, but this facility has never been granted. The wetland was also declared (a part of Ucchali Complex) a Ramsar Site on 22 March 1996. The surrounding agricultural land is privately owned whereas the hill forests and range lands are state owned (Government of the Punjab).
Current threats and problems	Illegal hunting, recreation, livestock grazing and agricultural activities cause some disturbance to waterbird, and the marked fluctuations in water level and salinity are having a detrimental effect on the habitat. There are reports of illegal hunting of waterbirds by local people and by military personnel stationed in the area. The Soil Conservation and Agriculture Departments advocates preventing run-off water from going into the lake by constructing bunds, and directing the water to the agricultural fields. In the process, land is reclaimed from the lake, which the owners use for cropping. The sinking of tube wells for irrigation purposes also affects the lake levels by drawing down the water table.
Suggestion for conservation of site	As for Jahlar Lake
Name of compiler	Abdul Aleem Chaudhry

Information Sheet 7. Important Sites for the White-headed Duck in Russia

Site Name	Kurlady Lake
Area (ha)	3,600 ha
Location	Southern Ural, Chelyabinsk Region, Kopeysk Town. 55°01'N, 61°05'E.
Site Description	The depth of the lake is about 3 m, it is weakly saline and has a silty bed. About 40% of the lake is covered by emergent vegetation (<i>Phragmites</i> , <i>Cyperacea</i> , <i>Scirpus</i>). The shoreline is covered by <i>Phragmites communis</i> .
Description and intensity of human use of this wetland	Fishing, agriculture, (animal grazing), hunting and recreation. There are buildings on the shore.
Protection/ ownership of site	Access to sections of the lake used by nesting birds is limited for people and animals due to the floating vegetation.
Current threats and problems	Fishing, disturbance from people, water level fluctuation
Suggestions for conservation of site	Roosting areas for the waterbirds need to be created close to the nesting sites.
Name of compiler	Gordienko N. S.

Site Name	Kulunda and Baraba forest steppes, Karasuksko-Burlaiskaya Lake System and Kulundinskiye Lakes
Area (ha)	There are four wetland complexes located within this large site (region). Their areas as defined for Ramsar designation are as follows: Chany Lakes 364,848 ha, Wetlands in the Lower Bagan area 26,880 ha, Karasuksko-Burlaiskaya Lake System more than 2,000 ha, and Kulundinskiye Lakes 200,000 ha. (Chany Lakes 55°16'30"N, 77°42'30"E and in Lower Bagan area 54°09'N, 78°23'E)
Location	South of Novosibirsk Region and west of Altai Territory
Site Description	Freshwater and saline lakes and adjacent land areas.
Description and intensity of human use of this wetland	Principal activities in and around the wetlands include agriculture, grazing, hay harvesting, hunting and fishing. In general, human pressure is higher in Chany and Lower Bagan, and lower in the two other wetland complexes.
Protection/ownership of site	Both Chany Lakes and Wetlands in the Lower Bagan Area are Ramsar sites, and both Karasuksko-Burlaiskaya Lake System and Kulundinskiye Lakes are on the Ramsar Shadow List. Many lakes, partly or fully, are included in Nature Reserves (Zakazniks). Lakes and adjacent lands are state (national) property or under collective proprietary rights by users.
Current threats and problems	Wetlands have a changeable regime of inundation and their areas can strongly decrease in some years.
Suggestions for conservation of site	It has been proposed to establish a strict nature reserve (Zapovednik) on Chany Lakes and another nature reserve (Zakaznik) in the Low Bagan area.
Name of compiler	Alexander Solokha, based on Krivenko 1999 and 2000.

Site Name	Lakes of Tobol-Ishim Forest-steppe
Area (ha)	1,217,000 ha (for the whole area of the wetland complex designated as a Ramsar site)
Location	55°00'–55°55'N, 67°05'–70°25'E. Ishim province of the forest-steppe zone on the Western Siberian Plain in the southern part of the Tyumen Region including territories of Armizonsky, Berdughsky, Kazansky, Sladkovsky and Ishimsky (partly) Administrative Districts, 190-250km south of the Tumen City.
Site Description	The wetland complex includes many lakes and marshes overgrown with emergent, floating and submerged aquatic plants (<i>Phragmites</i> , <i>Typha</i> , <i>Carex</i> , <i>Scirpus</i> , etc). The wetlands alternate with areas of birch and aspen forests as well as with meadows and steppes, most of which are ploughed.
Description and intensity of human use of this wetland	Agriculture, cattle grazing, fishing, waterbird hunting, recreation. All human activities affecting natural ecosystems are restricted (agriculture, fishing) or prohibited (hunting) in the protected areas.
Protection/ownership of site	Ramsar site. Most important wetland habitats are included in the 6 Nature Reserves and refuges comprising a total area of about 45,000 ha where hunting is prohibited and other human activities are limited. Most areas are under state ownership.
Current threats and problems	Waterbird hunting, especially in spring, impacts negatively on waterbird populations. For the last years, waterbird habitats have declined due to decreasing inundation of the forest-steppe.
Suggestions for conservation of site	It has been proposed to extend the total area of nature reserves by 30% and to include several lakes not yet protected. A number of measures aimed at limiting economic activities have been proposed, including restriction on grazing and fishing and prohibition of hunting during the breeding period.
Name of compiler	Alexander Solokha, based on Krivenko 1999.

Site Name	Ubsu-Nur Lake
Area (ha)	4490 ha
Location	50°40'N, 93°00'E. Republic of Tuva (south of Central Siberia) on the Mongolian border. In Mongolian side, the lake is called Uvs Lake.
Site Description	Saline lake with shoreline covered by reed communities and surrounded by mires and swamp meadows.
Description and intensity of human use of this wetland	Human use includes cattle grazing. Intensity is unclear but it is probably low.
Protection/ownership of site	This site belongs to the Ubsu-nur Depression Nature Reserve.
Current threats and problems	Unclear
Suggestions for conservation of site	None
Name of compiler	Alexander Solokha

Information Sheet 8. Important sites for the White-headed Duck in Turkmenistan

Site Name	Wetlands of the southeast coast of the Caspian Sea in Western Turkmenistan and filter reservoirs in Central and southeast Turkmenistan (combined sheet)
Area (ha)	More than 700,000 ha.
Location	Marine shallow coastal waters between cities Becdash, Crasnovodsk (now Turkmenbashies) and Gasankuly. Continental part of Turkmenistan (the area of the cities of Ashkhabad, Chardjou, Maryi and Tejen)
Site Description	The system of large marine embayments of the Caspian Sea - Rasnowodsky (now Turkmenbashies) and Severo-Chelekensky embayments are areas of Chasarskiy state natural reservation, the central and southeast Turkmenistan (Regions of the Caracum canal and Amu Darya River) artificial water storage basins, overflows, filtration lakes.
Description and intensity of human use of this wetland	Intensive economic and recreational activity, including hunting of waterbirds and fisheries.
Protection/ownership of site	Unclear
Current threats and problems	Overfishing, hunting, water allocation to inland wetlands
Suggestions for conservation of site	To undertake an ecological study of the areas and increase awareness of the importance of the area.
Name of compilers	M. E. Gauser and V. I. Vasilyev

Information Sheet 9. Important sites for the White-headed Duck in Uzbekistan

Site Name	Dengizkul Lake
Area (ha)	3,130 ha
Location	39°07'N, 64°10'E Altitude: 183.1 m ASL. This large saline water body in the southwest part of the Kyzylkum Desert lies in a natural depression 70km south-south-west of the city of Bukhara and 35km south-east of Alat Town on the border of Uzbekistan and Turkmenistan

Site Name	Dengizkul Lake
Site Description	<p>It is a lake with ecological conditions of natural lakes situated in the deserts of Central Asia. In the past, this lake was terminal the River Zaravshan. Records from 1914 indicate that the area of this lake was 120km². However, by the mid-1950s, the lake had completely dried up and turned into a big “solonchak” (big area of soil) because of the withdrawal of its waters for irrigation.</p> <p>Dengizkul Lake has no outflow. The main inflow is through the Dengizkul collector drain. With the increase in agriculture and water withdrawal, the drainage inflow into Dengizkul Lake grows.</p> <p>The construction of the Dengizkul collector drain and the Amu-Bukhara canal in 1966 resulted in an intensive filling of the lake. In the past 25 years, this lake has been converted into the biggest irrigation-wastewater lake from a periodically drying <i>solonchak</i>. The lake is 43.5km long aligned southeast to northwest. At its widest it is 9km with an average width of 7.2km.</p> <p>The coastline is stable. The depth in the central part is 23 m, with an average depth of about 10.69 m. The northeastern and southwestern reaches of the lake are separated from the central part by underwater elevations which impede water exchange between these areas and the central part.</p> <p>Most of the northeastern shore is steep, formed of soils with a high content of gypsum, sands and marl. The southern shore is low and covered with desert vegetation.</p> <p>Emergent vegetation is restricted to the northwestern part of the lake consisting mainly of reeds. About 4% of the lake area is covered with vegetation.</p> <p>The salinity of the lake at present is about 29.6mg/l. It increases as a result of evaporation.</p> <p>The climate is extreme continental, dry, with high fluctuations in temperature. An average temperature in January is 0 - 2⁰ C, the absolute minimum - 27⁰ C. The average temperature in July is +27-30⁰ C, but it can be as high as 45-48⁰C. During the winter, the lake freezes over for between 0 and 30 days. In cold winters the whole water surface of the lake may be frozen, but it usually covers only 20-30% of the surface area. This lake is of special value as a staging site for migrating waterbirds and as a wintering place for wetland birds.</p>
Description and intensity of human use of this wetland	<p>In 1992 Dengizkul Lake was developed for commercial fisheries. The fish yield on Dengizkul Lake reached 172 tonnes/year. At present, as a result of increasing water salinity this activity has ceased as fish have completely disappeared. The water level of the lake is decreasing each year by 0.5-1 m, and its bank zone has become shallow.</p> <p>Land adjoining this lake is used as desert pastures for livestock raising, mainly Astrakhan sheep. For the needs of the local population, small amounts of reed are harvested as construction material.</p> <p>The commercial mining of gas in the vicinity of and on Dengizkul Lake has enabled the provision of the local population with fuel and thus preserved trees and shrubs, which are important components of the desert ecosystem.</p>
Protection/ ownership of site	<p>Dengizkul Lake was recognised as a specially protected area or "Ornithological Zakaznik" in 1991 (category IV of IUCN). Dengizkul Lake was declared as a Ramsar site of Uzbekistan in February 2002. There are no special inspection points and staff at the lake. The lake is located at the frontier zone with Turkmenistan and it is not accessible without special permission.</p> <p>According to the Land Law of the Uzbekistan, the territory around Dengizkul Lake and the lake proper belongs to the state and is used by local people under control of the district and region khokimiyats - local authority, the Ministry of Agriculture and Water Management and Ministry of Oil and Gas Industry. State government bodies manage them on site.</p>
Current threats and problems	<p>Its hydrological regime depends on human activities. If water management authorities take the decision to stop drainage water into the lake, it is expected that there will be negative changes to the existing lake ecosystem. At present</p>

Site Name	Dengizkul Lake
	there is decreasing amount of water flowing in the lake from the Dengizkul collector. As a result, the mineral content of the water has increased to 29 g/l. It is expected that this will first lead to degradation of vegetation and reduction of fish, which will in time lead to a decline in numbers of fish-eating birds.
Suggestions for conservation of site	Undertake a public awareness campaign for local people, local authority and persons responsible for water management.
Name of compiler	Evgeniya Lanovenko

Site Name	Sudochoye Wetlands
Area (ha)	The Akushpa Lake area is about 3,070 ha
Location	58°30'N, 43°14'E Altitude: 52 m ASL. The Sudochoye Wetlands occupies a shallow and vast depression located in the northwest part of the Amu Darya River Delta (western part of the southern Aral Sea region). Nearest town was Muinak but at present, there are the towns of Kungrad (60km to the south-southeast) and the Roushan village (40km to the south).
Site Description	A large part of the wetland is occupied by large and small lakes and reed beds. The four main water bodies are: Akushpa Lake, Begdulla-Aydin Lake, Big Sudochoye Lake and Karateren Lake. Akushpa Lake is one the biggest among four large lakes, occupies the western part of the Sudochoye Wetlands adjoining the eastern edge of the "chink" Usturt Plateau. This is a saline lake. Its maximum length is about 20km and the maximum width is about 6.5km. The depth does not exceed 1.5-1.7 m. The total coastline length is about 62km. The wetland is surrounded with reed thickets, the height and density of which decreases towards its southern extremity. The open surface of the lake is about 50% of its total area. The water is very clear and transparent to the lake bottom. Hydrological and physico-chemical conditions of the water bodies depend on the relative amounts of inflowing and outflowing water. Mean duration of ice cover reaches 100 days, mainly restricted to the period of November 15 to March 10. Salinity of the Akushpa Lake fluctuates between 16.3 to 29.6 mg/l. In the summer and autumn of 2000, when the lake water began evaporating, the salinity rapidly increased.
Description and intensity of human use of this wetland	Local people use some sections of the Sudochoye Wetlands for fishing, animal grazing, hunting and catching of muskrats. In the past, people caught muskrats at the Akushpa Lake. At present both activities have been officially stopped and now a high level of poaching of muskrats occurs in autumn and early winter. During the breeding season of the birds there are no people at Akushpa Lake.
Protection/ ownership of site	The Sudochoye Wetlands is recognised as a specially protected area or "zakaznik" (category IV of IUCN). There are special inspection points and staff at the lake. According to the Law of the Republic of Uzbekistan on land, the territory around Sudochoye Wetlands and the lakes proper belong to the State and are used by the local people under the control of the district and region khokimiyats - local authority, the Ministry of Agriculture and Water Management. State bodies manage them on site.
Current threats and problems	The main current threat is the loss of habitats of the Sudochoye Wetlands as a result of the drying up of shallow lakes due to natural periods of drought in the Central Asian region. Due to the influence of those factors Sudochoye Wetlands totally dried out in 2001. Akushpa Lake was changed into a big swamp. It is possible that in future, natural climatic cycles will regularly affect the ecological conditions of the habitats (wetlands situated at the Aral Sea region) and therefore negatively affect White-headed Duck and others birds. Spring burning of reed beds is a widespread practice by local people.
Suggestions for	Sudochoye Wetlands should be nominated as a Ramsar site. It is of international

Site Name	Sudochye Wetlands
conservation of site	importance for conservation of Anatidae and other birds like grebes, pelicans, cormorants, herons, swans, eagles, falcons, shorebirds, gulls, terns, and passerines in spring, summer and autumn seasons. Awareness amongst the local people of the detrimental effect of their activities on the wetlands needs to be increased.
Name of compilers	Evgeniya Lanovenko and Elena Kreuzberg-Mukhina

Appendix C: Key to Scientific and English name of species referred in the report.

Birds

Scientific Name	English name
<i>Anas acuta</i>	Northern Pintail, Pintail
<i>Anas clypeata</i>	Shoveller
<i>Anas penelope</i>	European Wigeon
<i>Anas poecilorhyncha</i>	Spot-billed Duck
<i>Anas querquedula</i>	Garganey
<i>Anas strepera</i>	Gadwall
<i>Anser anser</i>	Greylag Goose
<i>Anser cygnoides</i>	Swan Goose
<i>Anser erythropus</i>	Lesser White-fronted Goose
<i>Anser indicus</i>	Bar-headed Goose
<i>Aquila heliaca</i>	Imperial Eagle
<i>Aythya ferina</i>	Common Pochard
<i>Aythya fuligula</i>	Tufted Duck
<i>Aythya nyroca</i>	Ferruginous Duck
<i>Branta ruficollis</i>	Red-breasted Geese
<i>Charadrius alexandrinus</i>	Kentish Plover
<i>Ciconia nigra</i>	Black Stork
<i>Circaetus gallicus</i>	Short-toed Eagle
<i>Dendrocygna javanica</i>	Lesser Whistling Duck
<i>Fulica atra</i>	Common Coot
<i>Grus grus</i>	Eurasian Crane
<i>Haliaeetus albicilla</i>	White-tailed Eagle
<i>Himantopus himantopus</i>	Black-winged Stilt

Scientific Name	English name
<i>Larus genei</i>	Slender-billed Gull
<i>Larus ridibundus</i>	Black-headed Gull
<i>Marmaronetta angustirostris</i>	Marbled Teal
<i>Mergellus albellus</i>	Smew
<i>Netta rufina</i>	Red-Crested Pochard
<i>Oxyura jamaicensis</i>	Ruddy Duck
<i>Oxyura leucocephala</i>	White-headed Duck
<i>Oxyura maccoa</i>	Maccoa Duck
<i>Pandion halietus</i>	Osprey
<i>Pelecanus crispus</i>	Dalmatian Pelican
<i>Pelecanus onocrotalus</i>	White Pelican
<i>Phalacrocorax pygmaeus</i>	Pygmy Cormorant
<i>Phoenicopterus ruber</i>	Greater Flamingo
<i>Platalea leucorodia</i>	Eurasian Spoonbill
<i>Plegadis falcinellus</i>	Glossy Ibis
<i>Podiceps cristata</i>	Great-crested Grebe
<i>Podiceps nigricollis</i>	Black-necked Grebe
<i>Remiz pendulinus</i>	Eurasian Penduline Tit
<i>Sarkidiornis melanotos</i>	Comb Duck
<i>Tachybaptus ruficollis</i>	Little Grebe
<i>Tadorna ferruginea</i>	Ruddy Shelduck
<i>Tadorna tadorna</i>	Common Shelduck

Plants

Scientific Name	English name
<i>Acacia modesta</i>	
<i>Adhatoda vasica</i>	Arusa
<i>Agropyron sp.</i>	Couch Grass
<i>Agropyron repens</i>	Couch Grass
<i>Alhagi camelorum</i>	Camel Thorn, Caspian Manna, Kameeldoringos
<i>Artemisia spp</i>	Sage Brush
<i>Asparagus gracilis</i>	Asparagus
<i>Atriplex cana</i>	Saltbush
<i>Bromus inermis</i>	Smooth Brome Grass
<i>Bromus sp.</i>	Brome Grass
<i>Calamagrostis epigeios</i>	Chee Reedgrass
<i>Carex fedia</i>	
<i>Chara sp</i>	Muskgrass, Stonewort
<i>Cocculus laeba</i>	
<i>Cynodon sp.</i>	Bermuda Grass
<i>Cynodon dactylon</i>	Bermuda Grass
<i>Cyperaceae sp.</i>	Sedge
<i>Cyperus eleusinoides</i>	Sedge
<i>Dactylis sp.</i>	Orchard Grass
<i>Ehretia laevis</i>	Chamaror

Scientific Name	English name
<i>Eichhornia crassipes</i>	Water Hyacinth
<i>Festuca sp.</i>	Fescue
<i>Festuca sulcata</i>	Fescue
<i>Gymnosporia royleana</i>	
<i>Halocnemum strobilaceum</i>	
<i>Haloxydon multiflorum</i>	Saksaul
<i>Hydrilla verticillata</i>	Florida Elodea
<i>Imperata cylindrica</i>	Cogon Grass
<i>Juncus sp.</i>	Rushes
<i>Melilotus indicus</i>	Yellow Sweetclover
<i>Najas sp.</i>	Naiad
<i>Najas marina</i>	Holly - leaved Naiad
<i>Nymphaea sp.</i>	Water Lily
<i>Olea ferruginea</i>	Elgon
<i>Paspalum distichum</i>	Water Finger Grass
<i>Phragmites sp.</i>	Common Reed
<i>Phragmites australis</i>	Common Reed
<i>Phragmites communis</i>	Common Reed
<i>Phyla nodiflora</i>	Turkey Tangle Fogfruit
<i>Potamogeton sp.</i>	Pondweed

Scientific Name	English name
<i>Potamogeton crispus</i>	Curly Pondweed
<i>Potamogeton pectinatus</i>	Fennel Pondweed
<i>Pyrethrum achilleifolium</i>	
<i>Ranunculus scleratus</i>	Celery-leaved Crowfoot
<i>Reptonia buxifolia</i>	Gurgura
<i>Rhazya stricta</i>	
<i>Rumex sp</i>	Dock
<i>Ruppia rostellata</i>	Ruppia
<i>Saccharum spontaneum</i>	Wild cane
<i>Sageretia lorandehuana</i>	
<i>Scirpus sp.</i>	Tule
<i>Scirpus sp. (short)</i>	Tule
<i>Scirpus sp. (tall)</i>	Tule
<i>Spergularia marina</i>	Salt-Marsh Sand Spurry

Scientific Name	English name
<i>Spiraea hypericifolia</i>	Iberian Spirea
<i>Stipa lessingiana</i>	
<i>Suaeda fruticosa</i>	Alkali Seepweed, Inkbush, Shrubby Seablite
<i>Tamarix aphyllia</i>	Eshel Haprakim
<i>Typha angustata</i>	Cat-tail
<i>Vallisneria spiralis</i>	Straight Vallis
<i>Withania coagulans</i>	Paneer Bandh
<i>Zizyphus nummularia</i>	Jujube, Ber
<i>Zannichellia palustris</i>	Horned Pondweed
<i>Zizyphus mauritiana</i>	Ber
<i>Zizyphus spp</i>	Jujube

Insects

Scientific Name	English name
<i>Corixa sp.</i>	Water Boatmen
<i>Micronecta sp.</i>	Water Boatmen

Fish

Scientific Name	English name
<i>Catla catla</i>	Catla
<i>Cirrhinus mrigala</i>	Mrigal
<i>Ctenopharyngodon idella</i>	Grass Carp
<i>Cyprinus carpio</i>	Common Carp
<i>Labeo rohita</i>	Indian Carp
<i>Oreochromis sp.</i>	Tilapia

Animals

Scientific Name	English name
<i>Canis aureus</i>	Golden Jackal
<i>Felis chaus</i>	Jungle Cat
<i>Felis libyca</i>	Libyan Wildcat
<i>Lepus nigricollis</i>	Indian Hare
<i>Phoca caspica</i>	Caspian Seal
<i>Sus scrofa</i>	Wild Boar
<i>Vulpes vulpes</i>	Red Fox
<i>Ondatra zibethicus</i>	Muskrat

Amphibians

Scientific Name	English name
<i>Rana tigrina</i>	Indian Bullfrog

Appendix D: Contact details for contributors to the national report questionnaire.

Overall range

Alexander Solokha
Wetlands International - Russia Office
Nikoloyamskaya Ulitsa, 19, strn.3, Moscow 109240
Russia
Phone: +7 095 7270939, Fax: +7 095 7270938
Asolokha@wwf.ru

Andy J. Green
Dept. of Conservation Biology, Doñana
Biological Station, Avda. María Luisa s/n, 41013
Sevilla, Spain
andy@ebd.csic.es

Baz Hughes
Head of Threatened Species
WWT, Slimbridge, Glos. GL2 7BT, UK
Baz.Hughes@wwt.org.uk

Joost van der Ven
PO Box 157, 3940 AD Doorn Netherlands
Javdv@mail.kg

China

He Fenqi
Institute Zoology Research, Chinese Academy of
Sciences, No. 19, Zhongguancun Road
Beijing 10080, P.R.China
cn_0707@sina.com

Ma Ming
Xinjiang Institute of Ecology and Geography
Chinese Academy of Sciences, No 40 Beijing
Road, Urumqi 830011, Xinjiang, P. R. China
Maming@ms.xjb.ac.cn

India

M. Zafar-ul Islam
Bombay Natural History Society
Hornbill House, Shaheed Bhagat Singh Road
Mumbai - 400 023, India
bnhs@bom4.vsnl.net.in

Iran

Behrouz Behrouzi-Rad
Faculty of Natural Resources and Marine Biology
Dept. of Environment, Noor City, I. R. Iran
Behrouzirad@Yahoo.com

Derek A. Scott
Ornithological Consultant
Castletownbere PO, Castletownbere,
County Cork, Ireland
Phone: +353 27 73327
Derekscott@eircom.net, derek_scott@talk21.com

Hamid Amini
Wildlife and Aquatic Affairs Bureau
Department of Environment
PO Box: 15875-5181, Tehran, I. R. Iran
Phone: +98 21 8269293; Fax: +98 21 8267993
Sadeghizadegan@abedi.net

Sadegh Sadeghi Zadegan
Wildlife and Aquatic Affairs Bureau
Department of Environment
PO Box: 15875-5181, Tehran, I. R. Iran
Phone: +98 21 8269293; Fax: +98 21 8267993
Sadeghizadegan@abedi.net

Yavar Shahbazi
Wildlife and Aquatic Affairs Bureau,
Department of Environment
PO Box: 15875-5181, Tehran, I. R. Iran
Phone: +98-21 8269293; Fax: +98-21 8267993
Sadeghizadegan@abedi.net

Kazakhstan

Andrew Grieve
Ornithological Society of The Middle East,
Caucasus and Central Asia, Rosemead, Grange
Road, Adlingfleet, Near Goole, East Yorkshire
DN14 8HZ, United Kingdom
Phone: +44 1724 798174, Fax: +44 1442 822623
Ag@osme.org

Lars Lachmann
NABU (German Society for Nature Conservation)
Project Office International, Invalidenstraße 112,
D-10115 Berlin, Germany
Phone: +49 30 284 984 44
Fax: +49 -30-284 984 84
Lars.Lachmann@NABU.de

Will Cresswell
Royal Society University Research Fellow
The Edward Grey Institute of Field Ornithology
Zoology Dept., South Parks Rd., Oxford, OX1 3PS
UK. Phone: +1865 271 203, +7773 800084
Willcresswell@supanet.com

Kyrgyz Republic

Alexander Yakovlev
Biosphere Reserve "Isyk-Kul"
Gagarin Street, 31, v. Ananyevo, Issyk-Kul region,
722324 Kyrgyz Republic
Phone: +996 3942 62194
yakovlev@ecology.elcat.kg

Mongolia

Axel Braunlich
Brüsseler Str. 46, D-13353 Berlin, Germany
Phone: +49-30-453 66 92
Braunlich@ipn.de

N. Tseveenmyadag
Researcher, Ornithological Laboratory
Institute of Biology, Academy of Sciences
Mongolia
Tseveenmyadag@magicnet.mn

Pakistan

Abdul Aleem Chaudhry
84-B-3, Johar Town, Lahore
Pakistan
mhaleemi@yahoo.com

Goetz Eichhorn
University of Groningen
Zoological Laboratory
PO Box 14, 9750 AA Haren
The Netherlands
eichgoetz@web.de

Sergey Yerokhov
MOH Institute of Zoology, Almaty 480032
Kazakhstan
instzoo@nurzat.kz

D. Salmakeyev
Biosphere Reserve "Isyk-Kul"
Gagarin Street, 31, v. Ananyevo, Issyk-Kul
region, 722324 Kyrgyz Republic
Phone: +996 3942 62194
issyk-kol@infotel.kg

Batdelger Dashnamjilyn
Mongolia Nature History Museum, Department of
Ornithology, Mongolian Bird Conservation Society
PO Box 120. Ulanbaatar 210136, Mongolia
Phone: +976-11-3245430, Fax: +976-11-323150
Birdbdr@yahoo.com

S.Gombobaatar
Lecturer, Zoology Department
Faculty of Biology, National University of
Mongolia, Ulaabaatar 210646A, PO Box 537,
Mongolia
Phone: +976-11-454809, Fax: +976-11-320159
Gomboo@www.com

Rahat Jabeen
WWF-Pakistan, Fortune Centre, Room # 606-607,
6th Floor, Block-6, P.E.C.H.S, Shahrah-i-Faisal,
Karachi 75400, Pakistan
Phone: +92-21 4544791-92 Fax: +92-21 4544790
rahatj69@hotmail.com

Pakistan

Zulfiqar Ali
Zoology Department Punjab, University Lahore,
Pakistan
Zulfiqarali68@yahoo.com

Russia

A. K. Yurlov
Institute of Systematic and Animal Ecology
Frunze str. 11, Novosibirsk-91, 630091, Russia
Phone: +7-383-2-170007, Fax: +7-383-2-170973
Ya@eco.nsc.ru, akyurlov@yahoo.com

Gordienko Nadejda Sergeevna
Ilmenski reserve 456317 Miass Chelybinsky
Province, Russia
Fax: +7-35135-7-02-86
Gordi@imeny.ac.ru

Tajikistan

Islom A. Abdusalyamov
Institute of Zoology and Parasitology
Academy of Sciences of Republic of Tajikistan
Home address: Rudaki Avenue, 85-57, Dushanbe,
734001 Tajikistan
Phone: +992 372 218154 (Home)
Islom@ac.tajik.net

Turkmenistan

Myrrhy Gauser
Ecological-enlightenment NGO “Southern-Caspian
Station”, Chasarskiy State natural reserve of the
Ministry research nature Turkmenistan.
19, Bahry Hasar str., Turkmenbashies 45000,
Turkmenistan
Phone: +8-993-24324116,
Fax +8-993-24324392 –for Vasilyev
Flamingo@cat.glasnet.ru, vpal_oil@rambler.ru

Vladislav Vasilyev
Ecological-enlightenment NGO “Southern-
Caspian Station”, Chasarskiy State natural reserve
of the Ministry research nature Turkmenistan.
19, Bahry Hasar str., Turkmenbashies 45000,
Turkmenistan
Phone: +8-993-24324116,
Fax: +8-993-24324392
Flamingo@cat.glasnet.ru, vpal_oil@rambler.ru

Uzbekistan

Alexander Filatov
Zoological Institute of Uzbek Academy of Sciences
Niyazov str. –1, 700095, Tashkent,
Republic of Uzbekistan
Filatov@comuz.uz

Elena Kreuzberg-Mukhina
Uzbekistan Zoological Society, Institute of
Zoology of Uzbek Academy of Sciences
Niyazov str. –1, 700095, Tashkent,
Republic of Uzbekistan
Phone: + 99871 1216185 Fax: + 99871 1206791
iucn_uz@mail.ru

Evgeniya Lanovenko
Uzbekistan Zoological Society
Zoological Institute of Uzbek Academy of Sciences
Niyazov str. –1, 700095, Tashkent,
Republic of Uzbekistan
Phone: +998-71-1445979
filatov@comuz.uz